5603HR

CHARACTERS IN DETAIL

Ear

Colour white Length short Density lax Shape-side profile tapering Awns or scurs - presence awns present Awns or scurs - distribution whole length Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium absent

Anthers - anthocyanin colouration

Lower glume

External surface smooth Extent of internal hairs strong - Group 3 Beak length medium to long moderately curved Beak curvature Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

slightly curved Beak curvature

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath weak

Flag leaf- blade (lower side) very weak to weak

Culm - neck weak

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration very strong erect to semi-erect Plant - growth habit

Time of ear emergence medium



AC BARRIE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)

Beak length short

Beak curvature v.slightly curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature straight to slightly curved

Grain

Colour red

Colouration with phenol very light to light

Straw

Length long to very long Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak to medium

Culm - neck medium
Ear medium

Plant growth characters

Flag leaf - anthocyanin of auricles

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit erect to semi-erect
Time of ear emergence very early to early



AMARETTO

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or Scurs - length: short to medium

Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3
Beak length: very short

Beak shape (curvature): slightly curved
Shoulder width narrow to medium

Shoulder shape straight

Lower Lemma

Beak shape: slightly to moderately curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: medium

Wall thickness @ 20cms below ear: medium to thick

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong
Culm neck: strong
Ear - Glaucosity: strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: intermediate Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



ASHBY

CHARACTERS IN DETAIL

Ear

Colour white Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length short to medium Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth Extent of internal hairs strong - Group 3 Beak length medium Beak curvature moderately curved

Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red Colouration with phenol medium

Straw

Length short to medium Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak Plant - growth habit intermediate Time of ear emergence late Flag leaf - anthocyanin of auricles very weak



AZURITE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface

Extent of internal hairs strong - Group 3

Beak length short

Beak curvature v.slightly curved

Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit semi-erect
Time of ear emergence medium



BELVOIR

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitylaxShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium Flag leaf - anthocyanin of auricles weak



BONZER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium
Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath very weak to weak
Flag leaf - blade (lower side) very weak to weak

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



CALIXO

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensityvery lax to laxShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length short
Beak curvature straight

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length long
Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath weak

Flag leaf- blade (lower side) very weak to weak

Culm - neck weak

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit semi-erect
Time of ear emergence early to medium



COBBER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3

Beak length short
Beak curvature straight
Shoulder width narrow
Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late



CORNETTO

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presenceawns presentAwns or scurs - distributionwhole lengthAwns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length long to very long
Beak curvature slightly curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature straight to slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length long

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) weak to medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit semi-erect
Time of ear emergence medium



DAFNE

CHARACTERS IN DETAIL

Ear

Colour: white

Length:medium to longDensity:lax to mediumShape (side profile)parallel sidedAwns or scurs - presence:scurs present

Awns or scurs - distribution:

Awns or Scurs - length: short to medium short to medium

Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: medium to strong
Beak length: very short to short

Beak shape (curvature): straight to slightly curved

Shoulder width medium to broad

Shoulder shape slightly sloping to straight

Lower Lemma

Beak shape:

Grain

Colour red Colouration with phenol: dark

Straw

Length: medium to long

Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong

Flag leaf - blade (lower side): medium to strong

Culm neck: strong
Ear - Glaucosity: strong

Plant growth characters

Coleoptile-anthocyanin weak

Plant - growth habit: semi-erect to intermediate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



EQUITY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthshort to medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong

Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour white

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit semi-erect
Time of ear emergence medium



ETHOS

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 medium to long Awns or scurs - length Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath weak to medium

Flag leaf- blade (lower side) weak

Culm - neck weak to medium

Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium



GRAFFITI

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium to long Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath very weak to weak
Flag leaf - blade (lower side) very weak to weak
Culm - neck very weak to weak
Ear weak

Plant growth characters

Coleoptile - anthocyanin colourationmediumPlant - growth habitsemi-erectTime of ear emergencemedium



GRANARY

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length long Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3

Beak length short to medium

Beak curvature slightly curved

Shoulder width narrow to medium

Shoulder shape sl. sloping to straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium
Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath weak to medium
Flag leaf - blade (lower side) weak to medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit semi-erect
Time of ear emergence medium
Flag leaf - anthocyanin of auricles



INFINITY

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth Extent of internal hairs weak (1) short to medium Beak length Beak curvature slightly curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

red Colouration with phenol dark

Straw

Length long to very long Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium Flag leaf- blade (lower side) weak Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong Plant - growth habit semi-erect

Time of ear emergence very early to early



KWS ALDERON

CHARACTERS IN DETAIL

Ear

Colour white
Length short
Density lax

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3

Beak length short

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short
Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak

Plant - growth habit semi-erect
Time of ear emergence medium



KWS BITTERN

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape sl. sloping to straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



KWS CHARING

CHARACTERS IN DETAIL

Ear

Colour white Length short Density lax

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong

Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour white

Colouration with phenol dark to very dark

Straw

Length medium Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath weak

Flag leaf- blade (lower side) very weak to weak
Culm - neck very weak to weak

Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit semi-erect
Time of ear emergence medium



SPRING WHEAT

AFP 1/2253

KWS COLLIER

CHARACTERS IN DETAIL

Ear

Colour white Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth Extent of internal hairs strong Beak length short

Beak curvature slightly curved Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour white

Colouration with phenol dark to very dark

Straw

short to medium Length Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak Plant - growth habit semi-erect Time of ear emergence early to medium



KWS FLINT

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surfacesmoothExtent of internal hairsstrong - Group 3Beak lengthshort to mediumBeak curvaturev.slightly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium



KWS KILBURN

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density very lax to lax Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length very long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KWS MULLINS

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length short to medium
Beak curvature v.slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red Colouration with phenol dark

Straw

Length very short Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



KWS RECOLETA

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long strong

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3

Beak length short

Beak curvature v.slightly curved Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

slightly curved Beak curvature

Grain

Colour

Colouration with phenol dark to very dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



Variety Name: KWS SCIROCCO Crop: SPRING WHEAT

Breeder's Reference: AFP No: 1/

Breeder: KWS LOCHOW GMBH Translated from:

Origin: GERMANY Date: 22 March 2013

Code: KWSC

Coleoptile-anthocyanin colouration: ABSENT TO VERY WEAK

*Plant - growth habit: SEMI-ERECT TO INTERMEDIATE

*Time ear emergence: EARLY TO MEDIUM

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: LOW

*Flag leaf - Glaucosity of sheath: WEAK TO MEDIUM
Flag leaf - Glaucosity of leaf blade: VERY WEAK TO WEAK

Anthers - anthocyanin colouration:

Glaucosity of neck: WEAK TO MEDIUM

Hairiness of uppermost node:

*Plant - height: MEDIUM TO LONG

*Straw - wall thickness @ 20cms below ear: VERY THIN *Ear - Glaucosity: WEAK

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: WHITE

*Ear - Shape (side profile):

Ear - Length:

*Ear - Density:

*Awns or scurs - presence:

PARELLEL SIDED

SHORT TO MEDIUM

LAX TO MEDIUM

SCURS PRESENT

*Awns or scurs - presence: Awns or scurs - distibution:

*Scurs at tip of ear - length: SHORT

*Awns at tip of ear - length: Lower glume - Beak length:

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: NARROW TO MEDIUM

Lower glume - Shoulder type: SLIGHTLY SLOPING TO STRAIGHT

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: WEAK

Lower glume - Internal imprint:

Lower lemma - Beak shape: SLIGHTLY CIRVED TO MODERATELY

Lower lemma - Beak length-: VERY SHORT TO SHORT

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: STRONG *Grain - colour: RED

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: LIGHT TO MEDIUM

*Seasonal type: SPRING

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015

SPRING WHEAT 2015

AFP 1/2354

KWS TALLAND

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long

Density lax
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length short to medium
Beak curvature v.slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit semi-erect
Time of ear emergence medium



KWS TORRIDON

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short to medium Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3 Beak length short to medium

slight to moderately curved Beak curvature

Shoulder width narrow to medium

elevated Shoulder shape

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium Plant - growth habit semi-erect Time of ear emergence early



KWS VALE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short to medium Apical rachis segment - convex face hairiness medium to strong

absent

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3 Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KWS WESTFIELD

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4 Awns or scurs - length long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth Extent of internal hairs strong - Group 3 Beak length short to medium slightly curved Beak curvature Shoulder width narrow

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

medium Length Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit semi-erect Time of ear emergence medium to late



KWS WILLOW

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short to medium Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3

Beak length short

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape sl. sloping to straight

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) weak to medium
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium



LENNOX

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium weak to medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3

Beak length short Beak curvature straight Shoulder width narrow Shoulder shape elevated

Lower lemma

slightly curved Beak curvature

Grain

Colour red Colouration with phenol very dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



LINTON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length short to medium
Beak curvature slightly curved
Shoulder width very narrow to narrow

Shoulder shape straight

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath weak
Flag leaf - blade (lower side) weak
Culm - neck weak
Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



MULIKA

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature moderately curved
Shoulder width negrow to medium

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath weak
Flag leaf- blade (lower side) weak
Culm - neck weak

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



Variety Name: MV KARIZMA Crop: SPRING WHEAT

Breeder's Reference: AFP No: 1/

Breeder: MTA MEZOGAZDASAGI **Translated from:**

Origin: HUNGREY Date: 22 March 2013

Code: KRZM

Coleoptile-anthocyanin colouration: ABSENT OR VERY WEAK

*Plant - growth habit: SEMI ERECT TO INTERMEDIATE

*Time ear emergence: EARLY

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: LOW TO MEDIUM

*Flag leaf - Glaucosity of sheath: STRONG
Flag leaf - Glaucosity of leaf blade: MEDIUM

Anthers - anthocyanin colouration:

Glaucosity of neck: STRONG

Hairiness of uppermost node:

*Plant - height: MEDIUM

*Straw - wall thickness @ 20cms below ear: THIN TO MEDIUM

*Ear - Glaucosity: WEAK

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: WHITE

*Ear - Shape (side profile): TAPERING
Ear - Length: LONG

*Ear - Density: VERY LAX TO LAX *Awns or scurs - presence: AWNS PRESENT

Awns or scurs - distibution: *Scurs at tip of ear - length:

*Awns at tip of ear - length:

Lower glume - Beak length:

MEDIUM

Lower glume - Beak shape (curvature): SLIGHTLY CURVED

Lower glume - Beak tip:

Lower glume - Shoulder width:NARROW TO MEDIUMLower glume - Shoulder type:STRAIGHT TO ELEVATED

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: WEAK

Lower glume - Internal imprint: Lower lemma - Beak shape:

Lower lemma - Beak length-:

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: MEDIUM *Grain - colour: RED

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

DARK

SPRING

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015

PARAGON

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length short

Beak curvature v.slightly curved Shoulder width medium

Shoulder shape sl. sloping to straight

Lower lemma

Beak curvature straight to slightly curved

Grain

Colour

Colouration with phenol medium to dark

Straw

medium Length Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side)

Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very strong Plant - growth habit semi-erect Time of ear emergence late to very late Flag leaf - anthocyanin of auricles absent or very weak



RAFFLES

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium Density lax to medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red Colouration with phenol dark

Straw

Length short
Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side)

Culm - neck weak to medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles very weak



RGT DOUBLESHOT

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong (3)
Beak length short

Beak curvature slightly curved Shoulder width medium to broad

Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol light

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colourationweak to mediumPlant - growth habitsemi-erectTime of ear emergencemedium to late



SLINKI

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium to dense

Shape-side profile parallel sided

Awas or sours a presence sours present

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong
Beak length medium

Beak curvature slight to moderately curved Shoulder width very narrow to narrow

Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

Colour white Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath weak
Flag leaf- blade (lower side) weak

Culm - neck weak to medium

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



Variety Name: SONETT Crop: SPRING WHEAT

Breeder's Reference: AFP No: 1

Breeder: LANTMANNEN SW SEED Translated from:

Origin: GERMANY Date: 16 April 2013

Code: SONE

Coleoptile-anthocyanin colouration: STRONG

*Plant - growth habit: SEMI-ERECT TO INTERMEDIATE

*Time ear emergence: EARLY TO MEDIUM

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath:

HIGH TO VERY HIGH

MEDIUM TO STRONG

Flag leaf - Glaucosity of leaf blade:

WEAK TO MEDIUM

Anthers - anthocyanin colouration:

Glaucosity of neck: MEDIUM

Hairiness of uppermost node:

*Plant - height: MEDIUM
*Straw - wall thickness @ 20cms below ear: VERY THIN

*Ear - Glaucosity: WEAK TO MEDIUM

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: WHITE
*Ear - Shape (side profile): TAPERING

Ear - Length: MEDIUM TO LONG

*Ear - Density: LAX

*Awns or scurs - presence: SCURS PRESENT

Awns or scurs - distibution:

*Scurs at tip of ear - length: SHORT TO MEDIUM

*Awns at tip of ear - length: Lower glume - Beak length:

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width:NARROW TO MEDIUMLower glume - Shoulder type:SLIGHTLY SLOPING

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: STRONG TO VERY STRONG

Lower glume - Internal imprint:

Lower lemma - Beak shape: STRAIGHT
Lower lemma - Beak length-: VERY SHORT

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour:

STRONG
RED

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: LIGHT TO MEDIUM

*Seasonal type: SPRING

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015

SPARROW

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long Apical rachis segment - convex face hairiness strong Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature straight

Grain

Colour

Colouration with phenol medium to dark

Straw

Length long Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath weak
Flag leaf- blade (lower side) weak
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration

Plant - growth habit semi-erect
Time of ear emergence medium



STRAVINSKY

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth Extent of internal hairs strong (3) Beak length medium

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol medium

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong Ear

strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



THURIO

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long

Density lax

Shape-side profile parallel sided
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length very long
Apical rachis segment - convex face hairiness medium

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs strong (3)

Beak length long to very long
Beak curvature moderately curved
Shoulder width absent or very narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium

Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit semi-erect
Time of ear emergence medium



TYBALT

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length short

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration

Lower glume

External surface

Extent of internal hairs strong - Group 3

Beak length short

Beak curvature slightly curved Shoulder width medium

Shoulder shape sl. sloping to straight

Lower lemma

Beak curvature straight

Grain

Colour red
Colouration with phenol medium

Straw

Length short
Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit semi-erect
Time of ear emergence medium to late



VARSITY

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long

Density lax

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong (3)
Pack langth

Beak length medium
Beak curvature moderate

Beak curvature moderately curved Shoulder width marrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium to strong medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit semi-erect
Time of ear emergence medium to late



ZIRCON

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature slightly curved Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration

Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



ZOLA

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong - Group 3
Beak length medium

Beak curvature strongly curved Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium

