

## WINTER BARLEY

AFP 2/2586

# ACUTE

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

#### Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	medium
Awn tips	weak

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

**AMARENA****CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium to dense
Length	medium to long
Shape	parallel

**Awn**

Length compared to ear	medium to long
------------------------	----------------

**Rachis**

Length of first segment	short
Curvature of first segment	weak

**Spikelet**

Sterile spikelet - attitude	
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	longer

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

## WINTER BARLEY

AFP 2/2447

# ARCHER

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax to medium
Length	medium
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	long
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	semi-prostrate to prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

**BAZOOKA****CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	short
Shape	parallel

**Awn**

Length compared to ear	long
------------------------	------

**Rachis**

Length of first segment	short to medium
Curvature of first segment	weak

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	weak
Lemma nerves - at early dough	very weak to weak
Awn tips	weak to medium

**Plant growth characters**

Plant - growth habit	intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

**BELFRY****CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	short
Shape	parallel

**Awn**

Length compared to ear	long
------------------------	------

**Rachis**

Length of first segment	short to medium
Curvature of first segment	weak

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

**Grain**

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

**Plant growth characters**

Plant - growth habit	intermediate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

**BLAZING****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	long
Shape	

**Awn**

Length compared to ear	medium
------------------------	--------

**Rachis**

Length of first segment	medium to long
Curvature of first segment	medium to strong

**Spikelet**

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/2581

# CADILLAC

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

#### Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	very weak to weak

#### Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/2448

# CALIFORNIA

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	long
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect



## WINTER BARLEY

AFP 2/1825

# CAMION

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax to medium
Length	long
Shape	tapering

#### Awn

Length compared to ear	+/- equal to longer
------------------------	---------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

#### Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	weak
Awn tips	weak

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	very early to early
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

## WINTER BARLEY

# CAMPANILE

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium to dense
Length	medium to long
Shape	tapering

#### Awn

Length compared to ear	short (shorter than ear)
------------------------	--------------------------

#### Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn cf grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium to strong
Rachilla hair type	short
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	horizontal to semi-recurved

## WINTER BARLEY

AFP 2/1606

# CARAT

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax to medium
Length	medium
Shape	tapering

#### Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

#### Rachis

Length of first segment	short to medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	horizontal to semi-recurved

## WINTER BARLEY

AFP 2/2058

# CASSATA

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium to dense
Length	medium
Shape	tapering

#### Awn

Length compared to ear	+/- equal to longer
------------------------	---------------------

#### Rachis

Length of first segment	short to medium
Curvature of first segment	weak to medium

#### Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	absent (grains thresh free)
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	medium to strong

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

## WINTER BARLEY

AFP 2/2582

# CAVALIER

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	medium to strong
Lemma nerves - at early dough	strong to very strong
Awn tips	strong to very strong

#### Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

**CELOONA****CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	dense
Length	short to medium
Shape	parallel

**Awn**

Length compared to ear	medium
------------------------	--------

**Rachis**

Length of first segment	short to medium
Curvature of first segment	weak

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

**Grain**

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	medium

**Plant growth characters**

Plant - growth habit	semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/1983

# CYPRESS

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	medium
Shape	parallel

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	medium
Lemma nerves - at early dough	strong
Awn tips	strong

#### Plant growth characters

Plant - growth habit	semi-prostrate to prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-recurved

Variety Name: **Daxor**

Crop: Winter Barley

**Breeders Reference:**

**Breeder:** Sarl Adrien Momont et Fils

**Origin:**

**Code:** DAXO

**AFP No:**

**Translated from:** French

**Date:** 24 March 2015

<b>*Plant - growth habit:</b>	Intermediate to semi prostrate
<b>*Lower leaves - hairiness of leaf sheaths:</b>	Absent
<b>*Time of ear emergence:</b>	
<b>*Flag leaf- anthocyanin colouration of auricles:</b>	Absent or very weak
<b>Frequency of plants with recurved flag leaves:</b>	Low
<b>Flag leaf- glaucosity of sheath:</b>	Medium to strong
<b>*Ear- glaucosity:</b>	Medium to strong
<b>Ear- attitude:</b>	Semi- erect
<b>Ear- length:</b>	Medium
<b>*Ear- number of rows:</b>	More than two
<b>*Ear- density:</b>	Medium to dense
<b>Ear- shape:</b>	Parallel
<b>*Awn- anthocyanin colouration of the tips:</b>	
<b>*Awn- length compared to ear:</b>	Medium to long
<b>Awn- spiculation of margins:</b>	
<b>*Sterile spikelet-attitude:</b>	
<b>Sterile spikelet-shape of tip:</b>	
<b>Sterile spikelet- length of lemma:</b>	
<b>Median spikelet- length of glume + awn of grain:</b>	Equal
<b>Lemma nerves- anthocyanin colouration:</b>	Absent to very weak
<b>*Plant height:</b>	
<b>Collar type:</b>	
<b>Rachis- length of first segment:</b>	Short to medium
<b>Rachis- width of first segment:</b>	
<b>Rachis- curvature of first segment:</b>	Weak
<b>Rachis- shape of first segment:</b>	
<b>Rachis(2-row)- humping of segments (mid third):</b>	
<b>Rachis(6-row)- degree of zig-zag (mid third):</b>	
<b>Kernel- colour of aleurone layer:</b>	Whitish
<b>*Grain- rachilla hair type:</b>	Long
<b>Grain- rachilla length:</b>	
<b>Grain- rachilla - length of tip hairs:</b>	
<b>Grain- spiculation of inner lateral nerves:</b>	Strong
<b>Grain- spiculation of outer lateral nerves:</b>	
<b>Grain- anthocyanin colouration of lemma nerves:</b>	Absent or very weak
<b>*Grain- ventral furrow - presence of hairs:</b>	Present
<b>*Grain- husk:</b>	Present
<b>Grain- length of lodicules ("collar" type only):</b>	
<b>Grain- disposition of lodicules:</b>	Clasping
<b>*Seasonal type:</b>	Winter
<b>Extra characters:</b>	

\* = Characteristics which should always be included in the description of the variety.

31 March 2015



**ELEMENT****CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	medium to long
Shape	parallel

**Awn**

Length compared to ear	medium
------------------------	--------

**Rachis**

Length of first segment	medium
Curvature of first segment	weak

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

**Grain**

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	very weak to weak
Lemma nerves - at early dough	weak to medium
Awn tips	very weak to weak

**Plant growth characters**

Plant - growth habit	semi-erect to intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

## WINTER BARLEY

AFP 2/2383

# ESCADRE

### CHARACTERS IN DETAIL

#### Ear

Number of rows	six
Density	dense
Length	short
Shape	parallel

#### Awn

Length compared to ear	+/- equal to longer
------------------------	---------------------

#### Rachis

Length of first segment	short to medium
Curvature of first segment	very weak

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	weak to medium
Awn tips	weak to medium

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

## WINTER BARLEY

AFP 2/2596

# FENTARA

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium to long
Curvature of first segment	weak

#### Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

Variety Name: **Findora**

Crop: Winter Barley

Breeders Reference:

AFP No: 2/2676

Breeder:

Translated from: German

Origin: Germany

Date: 09 April 2015

Code: FIND

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Early
*Flag leaf- anthocyanin colouration of auricles:	Absent or very week
Frequency of plants with recurved flag leaves:	Low
Flag leaf- glaucosity of sheath:	Medium
*Ear- glaucosity:	Medium
Ear- attitude:	Semi-erect to horizontal
Ear- length:	Medium
*Ear- number of rows:	Two
*Ear- density:	Dense
Ear- shape:	Tapering
*Awn- anthocyanin colouration of the tips:	Absent or very week
*Awn- length compared to ear:	Medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	N/A - non or rudimentary
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	Absent or very week
*Plant height:	Very short to short
Collar type:	
Rachis- length of first segment:	Short to medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Short
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Very week to week
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very week
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

\* = Characteristics which should always be included in the description of the variety.

09 April 2015

## WINTER BARLEY

AFP 2/1910

# FLAGON

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax to medium
Length	long
Shape	parallel

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	horizontal to semi-recurved

## WINTER BARLEY

AFP 2/2664

# FLETCHER

### CHARACTERS IN DETAIL

#### Ear

Number of rows	more than two
Density	dense
Length	short to medium
Shape	parallel

#### Awn

Length compared to ear	medium to long
------------------------	----------------

#### Rachis

Length of first segment	short to medium
Curvature of first segment	weak

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

#### Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	weak to medium

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/2392

# FLORENTINE

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	medium
Lemma nerves - at early dough	medium
Awn tips	medium

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect

**GALATION****CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	short to medium
Shape	parallel

**Awn**

Length compared to ear	medium
------------------------	--------

**Rachis**

Length of first segment	short to medium
Curvature of first segment	weak

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	weak to medium
Awn tips	weak

**Plant growth characters**

Plant - growth habit	semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	erect to semi-erect



**HARLEQUIN****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium
Shape	tapering

**Awn**

Length compared to ear	shorter to +/- equal
------------------------	----------------------

**Rachis**

Length of first segment	medium
Curvature of first segment	medium

**Spikelet**

Sterile spikelet - attitude	parallel
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

**Anthocyanin colouration**

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

**Plant growth characters**

Plant - growth habit	semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

**KATHMANDU****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

**Awn**

Length compared to ear	short
------------------------	-------

**Rachis**

Length of first segment	medium to long
Curvature of first segment	strong

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

**Plant growth characters**

Plant - growth habit	intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early
Flag leaf - glaucosity of sheath	weak
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

**KWS CAPELLA****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	very lax to lax
Length	short to medium
Shape	tapering

**Awn**

Length compared to ear	medium to long
------------------------	----------------

**Rachis**

Length of first segment	medium
Curvature of first segment	medium

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	very weak to weak

**Plant growth characters**

Plant - growth habit	semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/2309

## KWS CASSIA

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	medium
Shape	parallel

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	short
Curvature of first segment	weak to medium

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	very weak to weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

## WINTER BARLEY

AFP 2/2523

# KWS GLACIER

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	erect to semi-erect

**KWS INFINITY****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	very lax to lax
Length	short to medium
Shape	tapering

**Awn**

Length compared to ear	medium to long
------------------------	----------------

**Rachis**

Length of first segment	medium
Curvature of first segment	medium

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	very weak to weak

**Plant growth characters**

Plant - growth habit	semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/2521

# KWS JOY

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	long
Curvature of first segment	strong

#### Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium (3-5 per nerve)
Rachilla hair type	short
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/2436

# KWS MERIDIAN

### CHARACTERS IN DETAIL

#### Ear

Number of rows	six
Density	medium
Length	short to medium
Shape	parallel

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	strong

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect



**KWS ORWELL****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium
Shape	tapering

**Awn**

Length compared to ear	medium
------------------------	--------

**Rachis**

Length of first segment	long
Curvature of first segment	medium

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

**Plant growth characters**

Plant - growth habit	intermediate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	weak
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	erect to semi-erect

Variety Name: **KWS ROSE**

Crop: Winter Barley

Breeders Reference:

AFP No: 2/2730

Breeder:

Translated from:

Origin:

Date: 24 March 2015

Code: B2730

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Strong
Ear- attitude:	Semi - recurved to recurved
Ear- length:	Medium
*Ear- number of rows:	Two
*Ear- density:	Lax
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Very weak
*Awn- length compared to ear:	Short to medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	Parallel to weakly divergent
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	
Collar type:	
Rachis- length of first segment:	Medium to long
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

\* = Characteristics which should always be included in the description of the variety.

31 March 2015

**KWS SALSA****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium
Shape	tapering

**Awn**

Length compared to ear	medium to long
------------------------	----------------

**Rachis**

Length of first segment	medium to long
Curvature of first segment	weak to medium

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

**Plant growth characters**

Plant - growth habit	semi-erect to intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	weak to medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

**KWS TONIC****CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	dense
Length	medium
Shape	parallel

**Awn**

Length compared to ear	medium
------------------------	--------

**Rachis**

Length of first segment	short to medium
Curvature of first segment	very weak

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	very weak to weak
Lemma nerves - at early dough	weak to medium
Awn tips	medium

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

**KWS TOWER****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

**Awn**

Length compared to ear	medium
------------------------	--------

**Rachis**

Length of first segment	long
Curvature of first segment	strong

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	erect to semi-erect

**MARIS OTTER****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium to long
Shape	parallel

**Awn**

Length compared to ear	long (longer than ear)
------------------------	------------------------

**Rachis**

Length of first segment	medium to long
Curvature of first segment	weak to medium

**Spikelet**

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

**Anthocyanin colouration**

Flag leaf - auricles	strong
Lemma nerves - at early dough	medium
Awn tips	medium

**Plant growth characters**

Plant - growth habit	semi-prostrate to prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late to very late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	horizontal

## WINTER BARLEY

AFP 2/2452

# MATROS

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax to medium
Length	medium to long
Shape	parallel

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	short to medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	very strong
Lemma nerves - at early dough	strong to very strong
Awn tips	strong to very strong

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	strong
Ear - attitude 21 days after ear emerg.	semi-recurved to recurved

## WINTER BARLEY

AFP 2/2514

# MEZMAAR

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	short to medium
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect



## WINTER BARLEY

AFP 2/1318

## PEARL

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

#### Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn cf grain	equal

#### Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium (3-5 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	medium to strong
Lemma nerves - at early dough	very weak to weak
Awn tips	weak to medium

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	strong
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

## WINTER BARLEY

AFP 2/2651

# PERSEUS

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	short to medium
Shape	tapering

#### Awn

Length compared to ear	medium
------------------------	--------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

#### Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

#### Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	strong to very strong
Awn tips	strong

#### Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/2306

# PURDEY

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	medium to long
Shape	parallel

#### Awn

Length compared to ear	shorter to +/- equal
------------------------	----------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	semi-prostrate to prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

## WINTER BARLEY

AFP 2/2601

# QUADRA

### CHARACTERS IN DETAIL

#### Ear

Number of rows	more than two
Density	medium
Length	short to medium
Shape	parallel

#### Awn

Length compared to ear	medium
------------------------	--------

#### Rachis

Length of first segment	short to medium
Curvature of first segment	weak to medium

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	equal

#### Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

#### Anthocyanin colouration

Flag leaf - auricles	very weak to weak
Lemma nerves - at early dough	very weak to weak
Awn tips	weak

#### Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

**QUADRIGA****CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	dense
Length	short to medium
Shape	parallel

**Awn**

Length compared to ear	long
------------------------	------

**Rachis**

Length of first segment	medium
Curvature of first segment	medium

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	weak

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

**RETRIEVER****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium
Shape	tapering

**Awn**

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

**Rachis**

Length of first segment	medium
Curvature of first segment	medium

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium to strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

**Anthocyanin colouration**

Flag leaf - auricles	medium
Lemma nerves - at early dough	strong
Awn tips	strong

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

Variety Name: **Sabine F**

Crop: Winter Barley

Breeders Reference:

AFP No: 2/2231

Breeder: Saatzeit Filbeck

Translated from: German

Origin:

Date: 29 November 2007

Code: SABI

*Plant - growth habit:	intermediate
*Lower leaves - hairiness of leaf sheaths:	present
*Time of ear emergence:	late
*Flag leaf- anthocyanin colouration of auricles:	absent or very weak
Frequency of plants with recurved flag leaves:	high to very high
Flag leaf- glaucosity of sheath:	strong
*Ear- glaucosity:	weak to medium
Ear- attitude:	semi-erect to horizontal
Ear- length:	medium to long
*Ear- number of rows:	two
*Ear- density:	medium to dense
Ear- shape:	parallel
*Awn- anthocyanin colouration of the tips:	absent or very weak
*Awn- length compared to ear:	short to medium (shorter to +/- equal to ear)
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	parallel
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	medium
Collar type:	
Rachis- length of first segment:	long to very long
Rachis- width of first segment:	
Rachis- curvature of first segment:	weak to medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	whitish ("white")
*Grain- rachilla hair type:	long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	weak (1-2 per nerve)
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	absent or very weak
*Grain- ventral furrow - presence of hairs:	absent
*Grain- husk:	present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	clasping ("collar" type)
*Seasonal type:	winter
Extra characters:	

\* = Characteristics which should always be included in the description of the variety.

31 March 2015

## WINTER BARLEY

AFP 2/1880

# SAFFRON

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	medium
Shape	parallel

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	short to medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	absent or very weak
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal



## WINTER BARLEY

AFP 2/1717

# SEQUEL

### CHARACTERS IN DETAIL

#### Ear

Number of rows	six
Density	medium
Length	short to medium
Shape	tapering to parallel

#### Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

#### Rachis

Length of first segment	short
Curvature of first segment	weak

#### Spikelet

Sterile spikelet - attitude	
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	weak to medium

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

**SHADOW****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	short to medium
Shape	tapering

**Awn**

Length compared to ear	long
------------------------	------

**Rachis**

Length of first segment	medium to long
Curvature of first segment	strong

**Spikelet**

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	erect to semi-erect

## WINTER BARLEY

AFP 2/2454

# SINATRA

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/2513

# SOLOMAN

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	short
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

## WINTER BARLEY

AFP 2/2083

# SUZUKA

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering to parallel

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	short to medium
Curvature of first segment	medium

#### Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early
Flag leaf - glaucosity of sheath	weak to medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

## WINTER BARLEY

AFP 2/2444

## SY BAMBOO

### CHARACTERS IN DETAIL

#### Ear

Number of rows	more than two
Density	medium
Length	medium
Shape	parallel

#### Awn

Length compared to ear	medium
------------------------	--------

#### Rachis

Length of first segment	medium
Curvature of first segment	weak to medium

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	longer

#### Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

#### Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	weak to medium

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	long to very long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

**SY VENTURE****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium
Shape	tapering

**Awn**

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

**Rachis**

Length of first segment	medium
Curvature of first segment	medium to strong

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

**Anthocyanin colouration**

Flag leaf - auricles	strong
Lemma nerves - at early dough	very strong
Awn tips	strong

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

## WINTER BARLEY

AFP 2/2517

# TALISMAN

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

#### Awn

Length compared to ear	short (shorter than ear)
------------------------	--------------------------

#### Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	strong
Awn tips	strong

#### Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect



## WINTER BARLEY

AFP 2/2602

# TETRIS

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	lax
Length	long
Shape	tapering

#### Awn

Length compared to ear	short (shorter than ear)
------------------------	--------------------------

#### Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

#### Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	strong to very strong
Awn tips	strong

#### Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

**TRICK****CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium
Shape	parallel

**Awn**

Length compared to ear	medium
------------------------	--------

**Rachis**

Length of first segment	medium to long
Curvature of first segment	medium to strong

**Spikelet**

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

**Plant growth characters**

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

## WINTER BARLEY

AFP 2/2303

# WINSOME

### CHARACTERS IN DETAIL

#### Ear

Number of rows	two
Density	medium
Length	medium
Shape	parallel

#### Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

#### Rachis

Length of first segment	medium
Curvature of first segment	strong

#### Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

#### Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	short
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

#### Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	medium to strong
Awn tips	strong

#### Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

## WINTMALT

## CHARACTERS IN DETAIL

**Ear**

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering to parallel

**Awn**

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

**Rachis**

Length of first segment	medium
Curvature of first segment	strong

**Spikelet**

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	short
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

**Anthocyanin colouration**

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

Variety Name: **Verity**

Crop: Winter Barley

Breeders Reference:

AFP No: 2/2750

Breeder:

Translated from: German

Origin:

Date: 01 April 2015

Code: VERI

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Very Strong
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Semi erect
Ear- length:	Short
*Ear- number of rows:	More than two
*Ear- density:	Lax to medium
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	N/A - Deficiens
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Longer
Lemma nerves- anthocyanin colouration:	
*Plant height:	Long
Collar type:	
Rachis- length of first segment:	Medium to long
Rachis- width of first segment:	
Rachis- curvature of first segment:	Very weak to weak
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Medium to strong
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

\* = Characteristics which should always be included in the description of the variety.

09 April 2015

WINTER BARLEY 2015



14/04/2015 V1

**VOLUME****CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	medium
Shape	parallel

**Awn**

Length compared to ear	medium to long
------------------------	----------------

**Rachis**

Length of first segment	medium
Curvature of first segment	absent or very weak

**Spikelet**

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	equal

**Grain**

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

**Anthocyanin colouration**

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	very weak to weak
Awn tips	weak

**Plant growth characters**

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal