AEROBIC

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: dense

Shape (side profile) parallel sided Awns or scurs - presence: awns present

Awns or scurs - distribution:

Awns or Scurs - length: very long
Apical rachis segment - convex face hairiness: weak

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium - group 1 to 2

Beak length: medium to long

Beak shape (curvature): slightly curved

Shoulder narrow

Shoulder slightly sloping

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: medium

Straw

Length: short
Wall thickness @ 20cms below ear: very thick

Glaucosity

Flag leaf - sheath: weak to medium
Flag leaf - blade (lower side): very weak to weak

Culm neck: weak

Ear - Glaucosity: absent or very weak

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: semi-prostrate Time ear emergence: early

Flag leaf - Anthocyanin colouration of auricles:



ALCHEMY

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



ARARAT

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: lax
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: very weak to weak

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: medium - group 2

Beak length: medium

Beak shape (curvature): moderately curved

Shoulder medium Shoulder elevated

Lower Lemma

Beak shape: slightly curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: medium to strong
Flag leaf - blade (lower side): medium to strong

Culm neck: strong

Ear - Glaucosity: strong to very strong

Plant growth characters

Coleoptile-anthocyanin medium

Plant - growth habit: intermediate to semi-prostrate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



AVATAR

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitydenseShape-side profileparallel si

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit semi-erect
Time of ear emergence medium to late



BATTALION

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium - Group 2
Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence early to medium



BELENUS

CHARACTERS IN DETAIL

Ear

Colour: coloured

Length: short

Density: medium

Shape (side profile) parallel sided

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium
Apical rachis segment - convex face hairiness: weak

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak - group 1
Beak length: very short
Beak shape (curvature): slightly curved

Shoulder narrow

Shoulder slightly sloping

Lower Lemma

Beak shape: slightly curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: short
Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: weak

Flag leaf - blade (lower side): absent or very weak

Culm neck: absent or very weak

Ear - Glaucosity: absent or very weak

Plant growth characters

Coleoptile-anthocyanin absent or very weak

Plant - growth habit: intermediate
Time ear emergence: medium to late
Flag leaf - Anthocyanin colouration of auricles: absent or very weak



WINTER WHEAT

BELEPI

CHARACTERS IN DETAIL

Ear

Colour: white

Length: long to very long

Density: medium
Shape (side profile) parallel
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium
Apical rachis segment - convex face hairiness: weak

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length:

Beak shape (curvature):

Shoulder straight

Shoulder

Lower Lemma

Beak shape: straight to slightly curved

Grain

Colour red

Colouration with phenol: dark to very dark

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Plant growth characters

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): strong to very strong
Culm neck: strong to very strong

Ear - Glaucosity: strong to very strong

Coleoptile-anthocyanin weak

Plant - growth habit: intermediate to semi-prostrate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



BELUGA

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit semi-erect
Time of ear emergence medium



BISCAY

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-prostrate

Time of ear emergence late
Flag leaf - anthocyanin of auricles very weak



BOOMER

CHARACTERS IN DETAIL

Ear

Colour: white
Length: medium
Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong -group 3

Beak length: short

Beak shape (curvature): moderately curved

Shoulder narrow
Shoulder elevated

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: short
Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong
Culm neck: strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin strong
Plant - growth habit: semi-prostrate
Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



BROMPTON

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



BYSTRA

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense
Shape (side profile) parallel sided
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness: strong

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak - group 1
Beak length: very short

Beak shape (curvature): slightly curved

Shoulder narrow

Shoulder slightly sloping

Lower Lemma

Beak shape: slightly curved

Grain

Colour red

Colouration with phenol: light to medium

Straw

Length: short Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong
Culm neck: strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin weak
Plant - growth habit: semi-erect
Time ear emergence: medium to late
Flag leaf - Anthocyanin colouration of auricles: absent or very weak



CAMBRIDGE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 3/4
Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium



CASSIUS

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4

Awns or scurs - length: long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence late to very late



CHILTON

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution tip only Awns or scurs - length: long Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium - Group 2 Beak length medium to long Beak curvature moderately curved Shoulder width medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong

medium to strong Culm - neck

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong Plant - growth habit intermediate medium Time of ear emergence Flag leaf - anthocyanin of auricles

25/03/2014 V1

CHRONICLE

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature slightly curved
Shoulder width very narrow to narrow
Shoulder shape strongly elevated with 2nd pt.

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium



CLAIRE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: short

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Pook longth medium

Beak length medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



COCOON

CHARACTERS IN DETAIL

Ear

Colour white

Length medium

Density medium to dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs strong - Group 3

Beak length medium

Beak curvature moderately curved Shoulder width narrow shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) strong to very strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



CONQUEROR

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper 1/4Awns or scurs - length:short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



CONSORT

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper \(^{1}\)4
Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



CORDIALE

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length short

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape straight

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit medium to strong

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



CORONATION

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper 1/4

Awns or scurs - length: long
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong - Group 2+

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence medium to late



COUGAR

CHARACTERS IN DETAIL

Ear

Colour white Length short Density dense

Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: long to very long weak to medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium Beak curvature strongly curved Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



CRUSOE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width medium to broad

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath absent or very weak
Flag leaf- blade (lower side) absent or very weak
Culm - neck absent or very weak
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



CUBANITA

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: long

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late



DEBEN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



DELPHI

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 3/4
Awns or scurs - length: long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



DENMAN

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: short

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late Flag leaf - anthocyanin of auricles



DEXTER

absent

late

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium parallel sided Shape-side profile scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 medium to long Awns or scurs - length Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium

Beak length medium

Beak curvature strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence



DICKENS

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium



DUXFORD

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved

Shoulder width narrow

Shoulder shape strongly elevated with 2nd pt.

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) weak
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit semi-erect
Time of ear emergence early to medium



EDGAR

scurs present

CHARACTERS IN DETAIL

Ear

Colour: white
Length: long
Density: lax
Shape (side profile) tapering

Awns or scurs - presence:

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: weak to medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak - group 1
Beak length: very short to short
Beak shape (curvature): strongly curved

Shoulder narrow

Shoulder straight to elevated

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: medium Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong

Culm neck: strong to very strong
Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin very weak to weak Plant - growth habit: intermediate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



EINSTEIN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length she

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit semi-erect
Time of ear emergence early to medium
Flag leaf - anthocyanin of auricles absent or very weak



ENERGISE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence late

Time of ear emergence
Flag leaf - anthocyanin of auricles



EVOLUTION

CHARACTERS IN DETAIL

Ear

Colour white

Length medium

Density medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns of scurs - distribution upper ½

Avns or sours length.

Awns or scurs - length: long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



EXSEPT

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns of scurs - distributiontip onlyAwns or scurs - length:short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-prostrate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



FLORETT

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium - group 1 to 2

Beak length: very short to short
Beak shape (curvature): moderately curved

Shoulder narrow

Shoulder slightly sloping to straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: short
Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong

Flag leaf - blade (lower side): weak to medium

Culm neck: medium to strong

Ear - Glaucosity: medium

Plant growth characters

Coleoptile-anthocyanin absent or very weak
Plant - growth habit: intermediate
Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



GALLANT

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: medium
Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature straight to slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



GLADIATOR

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence early

Flag leaf - anthocyanin of auricles absent or very weak



GLASGOW

CHARACTERS IN DETAIL

Ear

Colour white

Length Short to medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence very early to early Flag leaf - anthocyanin of auricles absent or very weak



GOLDENGUN

CHARACTERS IN DETAIL

Ear

white Colour Length medium

Density medium to dense parallel sided Shape-side profile Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3 Beak length medium Beak curvature slightly curved Shoulder width narrow

Shoulder shape elev. to elev with 2nd point

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong to very strong Plant - growth habit intermed. to semi-prostrate

late

Time of ear emergence



GRAFTON

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium



GRAVITAS

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns of scurs - distribution upper ¹/₄
Awns or scurs - length: short

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



HAMPDEN

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 3/4 Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium Beak length medium to long Beak curvature moderately curved Shoulder width medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak intermediate Plant - growth habit

Time of ear emergence late



HAVANA

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit semi-erect
Time of ear emergence medium



HEREFORD

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or Scurs - length: long to very long

Apical rachis segment - convex face hairiness: strong

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length: medium

Beak shape (curvature): strongly curved

Shoulder medium
Shoulder elevated

Lower Lemma

Beak shape: strongly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong

Flag leaf - blade (lower side): strong to very strong

Culm neck: strong
Ear - Glaucosity: strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: intermediate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



HEREWARD

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns of scurs - distributionupper 1/4

Awns or scurs - length: very short to short

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium - Group 1+

Beak lengthmediumBeak curvatureslightly curvedShoulder widthbroad to very broad

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



HEROLDO

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: medium
Shape (side profile) tapering

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3
Beak length: short to medium

Beak shape (curvature): straight
Shoulder narrow

Shoulder straight to elevated

Lower Lemma

Beak shape:

Grain

Colour white

Colouration with phenol: light to medium

Straw

Length: short to medium

Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): strong to very strong

Culm neck: strong to very strong
Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: semi-prostrate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



HORATIO

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



HUMBER

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width medium to broad
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence early to medium



ICEBREAKER

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium
Beak curvature slightly curved
Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late



ICON

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long Density medium Shape-side profile tapering Awns or scurs - presence scurs present

Awns of scurs - distribution upper 3/4 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium to long Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Straw

Colour red Colouration with phenol dark

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit

intermed. to semi-prostrate

Time of ear emergence medium to late Flag leaf - anthocyanin of auricles



INVICTA

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributiontip onlyAwns or scurs - length:short to medium

Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late



ISTABRAQ

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper 1/4Awns or scurs - length:medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect

Time of ear emergence very early to early

Flag leaf - anthocyanin of auricles absent or very weak



JB DIEGO

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution tip only

Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length

Beak curvature slight to moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong weak to medium

Ear

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



JORVIK

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape straight

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



KETCHUM

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness medium

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red Colouration with phenol dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KINGDOM

absent

CHARACTERS IN DETAIL

Anthers - anthocyanin colouration

Ear

white Colour Length medium Density medium to dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution tip only Awns or scurs - length: short Apical rachis segment - convex face hairiness medium

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



KIPLING

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitydense to very denseShape-side profileparallel sided

Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: short to medium
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side)

Culm - neck medium Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KRIS

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium

Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3
Beak length: short to medium
Beak shape (curvature): slightly curved

Shoulder medium

Shoulder straight to elevated

Lower Lemma

Beak shape: slightly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: short
Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): medium to strong

Culm neck: strong
Ear - Glaucosity: strong

Plant growth characters

Coleoptile-anthocyanin medium to strong

Plant - growth habit: semi-prostrate to prostrate

Time ear emergence: late

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



KWS BONHAM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper 1/4Awns or scurs - length:medium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width marrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium
Culm - neck medium

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS CASHEL

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length very long
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong

Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath weak

Flag leaf- blade (lower side) very weak to weak

Culm - neck weak Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

late

Time of ear emergence



KWS CHILOE

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence awns present Awns of scurs - distribution whole length Awns or scurs - length: very long Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong - Group 3
Beak length long to very long

Beak curvature slightly curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol very light to light

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



KWS CLEVELAND

CHARACTERS IN DETAIL

Ear

Colour white
Length short to medium
Density medium

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper \(^{1}\)4
Awns or scurs - length: long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit semi-erect
Time of ear emergence medium



KWS CROFT

weak

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3 Beak length medium Beak curvature slightly curved Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour very dark

Colouration with phenol

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



KWS CROSBY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: long weak

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2 Beak length medium to long Beak curvature moderately curved narrow to medium Shoulder width

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong

Culm - neck strong to very strong Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KWS DALI

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long

Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS EVOKE

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness absent to weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red Colouration with phenol dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong Plant - growth habit intermediate medium Time of ear emergence



KWS EVOLVE

CHARACTERS IN DETAIL

Ear

Colour white Length long Density medium parallel sided Shape-side profile scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 medium Awns or scurs - length Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium
Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence late



KWS GATOR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS HORIZON

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness absent to weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitsemi-erectTime of ear emergencemedium to late



KWS KENTON

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved narrow to medium Shoulder width

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) weak to medium Culm - neck weak to medium Ear

medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong Plant - growth habit intermediate Time of ear emergence medium to late



KWS KIELDER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns of scurs - distributionupper ½Awns or scurs - length:long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature slightly curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit semi-prostrate
Time of ear emergence medium to late



KWS LILI

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium
Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium

Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



KWS LUMOS

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium Density medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium Beak curvature strongly curved Shoulder width narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red Colouration with phenol medium

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong Plant - growth habit semi-erect Time of ear emergence medium



KWS MADRYN

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: medium weak to medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium Beak curvature strongly curved Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



KWS MOJITO

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution upper 3/4
Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium

Beak length medium

Beak curvature moderately curved Shoulder width marrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak to medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence late

Time of ear emergence
Flag leaf - anthocyanin of auricles



KWS MOZART

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium parallel sided Shape-side profile scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/2 short to medium Awns or scurs - length Apical rachis segment - convex face hairiness medium

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



KWS PODIUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red Colouration with phenol dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit semi-erect
Time of ear emergence medium



KWS ROWAN

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium



KWS SANTIAGO

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KWS SAXTEAD

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence awns present Awns of scurs - distribution whole length Awns or scurs - length: long Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs strong - Group 3

Beak length long

Beak curvature strongly curved Shoulder width narrow elevated

Shoulder shape

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong

Culm - neck strong to very strong Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong Plant - growth habit semi-erect Time of ear emergence medium



KWS SOLO

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper 1/4

Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS STERLING

CHARACTERS IN DETAIL

Ear

Colour white

Length
Density
Shape-side profile
Awns or scurs - presence
Awns of scurs - distribution
Awns or scurs - length:
Apical rachis segment - convex face hairiness

short to medium
parallel sided
scurs present
upper \(^1\)4
long
weak

Anthers - anthocyanin colouration weak absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



KWS TARGET

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: medium
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitsemi-erectTime of ear emergencemedium



KWS TEMPO

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium
Beak length short to medium
Beak curvature strongly curved
Shoulder width narrow

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence very late



KWS TRINITY

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 medium Awns or scurs - length Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium
Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence late

Time of ear emergence
Flag leaf - anthocyanin of auricles



KWS YARIS

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium

Apical rachis segment - convex face hairiness: medium to strong

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length: long

Beak shape (curvature): strongly curved

Shoulder medium
Shoulder elevated

Lower Lemma

Beak shape: strongly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: medium Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong to very strong

Flag leaf - blade (lower side): very strong

Culm neck: strong to very strong
Ear - Glaucosity: strong to very strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: semi-prostrate

Time ear emergence: late

Flag leaf - Anthocyanin colouration of auricles:



LANCASTER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - length:long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width marrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck weak to medium
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

late

Time of ear emergence



LEAR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



LEEDS

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper ½Awns or scurs - length:long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium



MAGISTER

CHARACTERS IN DETAIL

Ear

Colour: white
Length: medium
Density: lax to medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium -group 1 to 2

Beak length: short

Beak shape (curvature): moderately curved
Shoulder medium to broad

Shoulder straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: medium to long Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong
Culm neck: strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin weak

Plant - growth habit: semi-prostrate
Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



MARIS WIDGEON

CHARACTERS IN DETAIL

white Colour Length medium Density very lax to lax tapering Shape-side profile Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs weak - Group 1
Beak length medium
Beak curvature v.slightly curved
Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very long Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence late
Flag leaf - anthocyanin of auricles weak



MONTEREY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



MYRIAD

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¾

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



NELSON

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



OAKLEY

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium



OLIVIN

CHARACTERS IN DETAIL

Ear

Colour: white
Length: medium
Density: medium
Shape (side profile) parallel sided
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: medium to strong

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: medium - group 2
Beak length: very short to short
Beak shape (curvature): strongly curved
Shoulder narrow to medium

Shoulder slightly sloping to straight

Lower Lemma

Beak shape: strongly curved to geniculate

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: medium to long

Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): medium
Culm neck: medium

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin strong to very strong

Plant - growth habit: intermediate to semi-prostrate

Time ear emergence: late

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



PANACEA

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¾

Awns or scurs - length: long
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

late

Time of ear emergence



PANORAMA

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length: short to medium
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium
Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence 1



PHILOSOPHER

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length: long
Apical rachis segment - convex face hairiness weak

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



PIRENEO

CHARACTERS IN DETAIL

Ear

Colour: white Length: medium

Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: awns present

Awns or scurs - distribution:

Awns or scurs - length: long
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: medium - group 2
Beak length: long to very long
Beak shape (curvature): slightly curved

Shoulder narrow Shoulder straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: medium

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: weak

Flag leaf - blade (lower side): very weak to weak

Culm neck: weak to medium

Ear - Glaucosity: weak

Plant growth characters

Coleoptile-anthocyanin

Plant - growth habit: intermediate
Time ear emergence: early to late

Flag leaf - Anthocyanin colouration of auricles:



QI

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - length:medium

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit semi-erect
Time of ear emergence medium



RA3W 08

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness: weak to medium

Anthers - anthocyanin

Lower Glume

External surface roughness: medium

Extent of internal hairs: medium - group 2

Beak length: medium

Beak shape (curvature): moderately curved

Shoulder broad
Shoulder elevated

Lower Lemma

Beak shape: moderately curved to strongly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: short to medium

Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): medium to strong

Culm neck: weak
Ear - Glaucosity: medium

Plant growth characters

Coleoptile-anthocyanin medium
Plant - growth habit: intermediate
Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



RAPSODIA

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium

Density: medium

Shape (side profile) tapering

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: weak to medium

Anthers - anthocyanin

Lower Glume

External surface roughness: smooth

Extent of internal hairs: medium to strong - group 2 to 3

Beak length: short

Beak shape (curvature): moderately curved

Shoulder medium Shoulder straight

Lower Lemma

Beak shape: slightly curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: short

Wall thickness @ 20cms below ear: very thin to thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): medium to strong

Culm neck: medium to strong
Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin strong

Plant - growth habit: semi-prostrate to prostrate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



RAZZANO

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length: very short
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface
Extent of internal hairs medium to strong - Group 2+

Beak length short

Beak curvature v.slightly curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



REFLECTION

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthmediumApical rachis segment - convex face hairinessabsent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium to late



RELAY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ½

Awns or scurs - length: long
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



REVELATION

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length: long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium - Group 2 Beak length medium

Beak curvature moderately curved Shoulder width medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour Colouration with phenol very dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



RGT CONVERSION

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length long

Beak curvature strongly curved Shoulder width narrow

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence late



ROBIGUS

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium
Flag leaf - anthocyanin of auricles weak



RUSKIN

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium
Beak length short to medium
Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



RW41138

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium to late



SAHARA

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ½

Awns or scurs - distribution upper 72

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak

Culm - neck weak

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



SCOUT

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature straight to slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



SCRIBE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: medium
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

anthory unithocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence medium



SKYFALL

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence awns present Awns or scurs - distribution whole length Awns or scurs - length: very long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong - Group 3
Beak length long to very long
Beak curvature v.slightly curved
Shoulder width very narrow to narrow
Shoulder shape elev. to elev with 2nd point

Lower lemma

Beak curvature straight

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium



SOISSONS

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length:

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak - Group 1

Beak length

Beak curvature

Shoulder width narrow

Shoulder shape slightly sloping

Lower lemma

Beak curvature straight

Grain

Colour red

Colouration with phenol

Straw

Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath absent or very weak

Flag leaf- blade (lower side) weak

Culm - neck absent or very weak Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence very early to early Flag leaf - anthocyanin of auricles absent or very weak



SOLACE

CHARACTERS IN DETAIL

Ear

Colour white

Length
Density
Shape-side profile
Awns or scurs - presence
Awns or scurs - distribution
Awns or scurs - length:
Apical rachis segment - convex face hairiness

short to medium
parallel sided
scurs present
upper ½
long
medium

Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



SOLSTICE

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence late
Flag leaf - anthocyanin of auricles very weak



STIGG

short to medium

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium Density very dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration absent

Lower glume

Awns or scurs - length:

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

moderately curved Beak curvature Shoulder width narrow to medium Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour

Colouration with phenol medium to dark

Straw

Length short thin

Wall thickness @20cms below ear

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



SY EPSON

CHARACTERS IN DETAIL

Ear

Colour: white Length: long

Density: lax to medium
Shape (side profile) parallel sided
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: long
Apical rachis segment - convex face hairiness: very weak

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium - group 1 to 2

Beak length: short to medium

Beak shape (curvature): slightly to moderately curved

Shoulder narrow to medium

Shoulder slightly sloping or straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: medium

Straw

Length:

Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: medium to strong

Flag leaf - blade (lower side): medium

Culm neck: medium to strong
Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin absent to very weak Plant - growth habit: semi-prostrate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



TIMARU

CHARACTERS IN DETAIL

Ear

Colour white Length long Density dense

Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution tip only Awns or scurs - length: very short Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium Beak curvature strongly curved narrow to medium Shoulder width

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong medium to strong

Ear

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate late

Time of ear emergence Flag leaf - anthocyanin of auricles

25/03/2014 V1

TORCH

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - length:medium

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



TORPHINS

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium to late



TUXEDO

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: medium to long

Aprical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



TWISTER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ½

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium



VISCOUNT

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



WARRIOR

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear n/a

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



WEAVER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length: long
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium

Beak curvature strongly curved Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence medium



WELFORD

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong

weak to medium Ear

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



XI19

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long

Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - length: short

Awns or scurs - distribution tip only
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late
Flag leaf - anthocyanin of auricles very weak



ZEBEDEE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



ZULU

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4 Awns or scurs - length: long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak to medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late

