Variety Name: **Aarden** Crop: Winter Wheat

Breeder's Reference: SUR 97271/10 **AFP No:** 1/1573

Breeder: Saaten Union Recherche SARL Translated from: German

Origin: Date: 19 November 2003

Code: ARDN

Coleoptile-anthocyanin colouration: medium

*Plant - growth habit: semi-prostrate to prostrate

*Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: absent or very low

*Flag leaf - Glaucosity of sheath: strong
Flag leaf - Glaucosity of leaf blade: strong
Anthers - anthocyanin colouration:

Glaucosity of neck: strong to very strong

Hairiness of uppermost node:

*Plant - height: short

*Straw - wall thickness @ 20cms below ear: very thin

*Ear - Glaucosity: strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: white

*Ear - Shape (side profile): tapering
Ear - Length: medium

*Ear - Density: medium

*Awns or scurs - presence: scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: short

*Awns at tip of ear - length: Lower glume - Beak length:

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: Lower glume - Shoulder type:

Lower glume - External surface roughness: Lower glume - Extent of internal hairs: Lower glume - Internal imprint:

Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour: red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: medium *Seasonal type: winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 06 May 2020

AEROBIC

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: dense

Shape (side profile) parallel sided Awns or scurs - presence: awns present

Awns or scurs - distribution:

Awns or Scurs - length: very long
Apical rachis segment - convex face hairiness: weak

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium - group 1 to 2

Beak length: medium to long
Beak shape (curvature): slightly curved

Shoulder narrow

Shoulder slightly sloping

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: medium

Straw

Length: short
Wall thickness @ 20cms below ear: very thick

Glaucosity

Flag leaf - sheath: weak to medium
Flag leaf - blade (lower side): very weak to weak

Culm neck: weak

Ear - Glaucosity: absent or very weak

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: semi-prostrate Time ear emergence: early

Flag leaf - Anthocyanin colouration of auricles:



ALCHEMY

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)
Culm - neck

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



AMBOISE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence medium



AMBROSIA

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



AMPLIFY

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong (2+)
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



Variety Name: **ANAPOLIS** Crop: Winter wheat

Breeder's Reference: AFP No: 1/2148

Breeder: NORDSAAT Translated from: German

Origin: Date: 23 July 2013

Code: ANAP

Coleoptile-anthocyanin colouration:Medium to strong*Plant - growth habit:Semi prostrate*Time ear emergence:Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath:

Very low to low

Strong to very strong

Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Absent or very thin.

*Ear - Glaucosity: Strong to very strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Tapering

Ear - Length: Medium

*Ear - Density: Lax to medium

*Awns or scurs - presence: Present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length:Very short to short **Lower glume - Beak shape (curvature):**Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width:Narrow **Lower glume - Shoulder type:**Straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Medium

Lower glume - Internal imprint:

Lower lemma - Beak shape: Slightly curved

Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour:

Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Medium to dark

*Seasonal type: Winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety.

ARARAT

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: lax
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: very weak to weak

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: medium - group 2

Beak length: medium

Beak shape (curvature): moderately curved

Shoulder medium
Shoulder elevated

Lower Lemma

Beak shape: slightly curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: medium to strong
Flag leaf - blade (lower side): medium to strong

Culm neck: strong

Ear - Glaucosity: strong to very strong

Plant growth characters

Coleoptile-anthocyanin medium

Plant - growth habit: intermediate to semi-prostrate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



Variety Name: Arkeos

Crop:

Breeder's Reference:

AFP No:

Breeder: Nickerson International Research

Translated from:

Origin:

Date:

26 January 2015

Code: ARKE

Coleoptile-anthocyanin colouration:

Medium

*Plant - growth habit:

Semi- prostrate to prostrate

*Time ear emergence:

Early

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath: Flag leaf - Glaucosity of leaf blade: Anthers - anthocyanin colouration: Strong to very strong Strong to very strong

Glaucosity of neck:

Hairiness of uppermost node:

*Plant - height: Short to medium
*Straw - wall thickness @ 20cms below ear: Very thin
*Ear - Glaucosity: Medium

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile):

Ear - Length: Long

*Ear - Density: Lax to medium

*Awns or scurs - presence: Present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Short to medium

*Awns at tip of ear - length:

Lower glume - Beak length: Short to medium

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow

Lower glume - Shoulder type: Straight to slightly sloping

Lower glume - External surface roughness: Lower glume - Extent of internal hairs:

Lower glume - Internal imprint:Medium to strongLower lemma - Beak shape:Slightly curvedLower lemma - Beak length-:Short to medium

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

Mdium to dark
Winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015

ASTOUND

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long

Density dense

Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs medium to strong (group 2+)

Beak length long

Beak curvature strongly curved Shoulder width medium

Shoulder shape elev. to strongly elevated

Lower lemma

Beak curvature geniculate

Grain

Colouration with phenol very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence

Flag leaf - anthocyanin of auricles absent or very weak



AVATAR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density dense
Shape-side profile parallel s

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit semi-erect
Time of ear emergence medium to late



Variety Name: Axioma Crop: Wheat

Breeder's Reference: SEC 228-02-4 **AFP No:**

Breeder: Secobra Rechereches Translated from: German

Origin: Date: 19 November 2018

Code: Axio

Coleoptile-anthocyanin colouration:Medium*Plant - growth habit:Intermediate*Time ear emergence:Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Very low to low

*Flag leaf - Glaucosity of sheath: Srong
Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Medium *Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour:

*Ear - Shape (side profile): Tapering
Ear - Length: Short to medium

*Ear - Density: Dense *Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature): Moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow to medium

Lower glume - Shoulder type: Straight

Lower glume - External surface roughness: Lower glume - Extent of internal hairs: Lower glume - Internal imprint: Lower lemma - Beak shape:

Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

Very dark
Winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 20 March 2019

BAHAMAS CS

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



BANQUO

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf - blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence lat

Flag leaf - anthocyanin of auricles absent or very weak



BARKSTON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length shor

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) medium to strong

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence very late



BATSMAN

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthshort to medium

Apical rachis segment - convex face hairiness
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence late to very late



BATTALION

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium - Group 2
Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence early to medium



BELENUS

CHARACTERS IN DETAIL

Ear

Colour:colouredLength:shortDensity:mediumShape (side profile)parallel sidedAwns or scurs - presence:scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium
Apical rachis segment - convex face hairiness: weak

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak - group 1
Beak length: very short
Beak shape (curvature): slightly curved

Shoulder narrow

Shoulder slightly sloping

Lower Lemma

Beak shape: slightly curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: short
Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: weak

Flag leaf - blade (lower side):

Culm neck:

absent or very weak

Ear - Glaucosity:

absent or very weak

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: intermediate Time ear emergence: medium to late Flag leaf - Anthocyanin colouration of auricles: absent or very weak



WINTER WHEAT

BELEPI

CHARACTERS IN DETAIL

Ear

Colour: white

Length: long to very long

Density: medium
Shape (side profile) parallel
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium
Apical rachis segment - convex face hairiness: weak

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length:

Beak shape (curvature):

Shoulder straight

Shoulder

Lower Lemma

Beak shape: straight to slightly curved

Grain

Colour red

Colouration with phenol: dark to very dark

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): strong to very strong
Culm neck: strong to very strong
Ear - Glaucosity: strong to very strong

Plant growth characters

Coleoptile-anthocyanin weak

Plant - growth habit: intermediate to semi-prostrate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



BELGRADE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length long to very long

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface rough
Extent of internal hairs strong (3)
Beak length medium to long
Beak curvature moderately curved
Shoulder width

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - peck medium

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



BELUGA

CHARACTERS IN DETAIL

Ear

Colour white Length medium lax to medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colourationweakPlant - growth habitsemi-erectTime of ear emergencemedium



BENNINGTON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlongApical rachis segment - convex face hairinessstrong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence very late



BISCAY

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-prostrate

Time of ear emergence late
Flag leaf - anthocyanin of auricles very weak



BLACKSMITH

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness absent or very small

Anthers - anthocyanin colouration present

Lower glume

External surface smooth
Extent of internal hairs weak (group 1)
Beak length short to medium
Beak curvature slightly curved
Shoulder width very narrow to narro

Shoulder width very narrow to narrow Shoulder shape slightly elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour purple
Colouration with phenol very dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



BOOMER

CHARACTERS IN DETAIL

Ear

Colour: white
Length: medium
Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong -group 3

Beak length: short

Beak shape (curvature): moderately curved

Shoulder narrow
Shoulder elevated

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: short
Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong
Culm neck: strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyaninstrongPlant - growth habit:semi-prostrateTime ear emergence:medium

Flag leaf - Anthocyanin colouration of auricles:



BRIDGEHEAD

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colourationweakPlant - growth habitintermediateTime of ear emergencelate to very late



BRITANNIA

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long

Density lax

Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 3/4 Awns or scurs - length very long Apical rachis segment - convex face hairiness strong absent

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs weak to medium (1+) Beak length medium to long Beak curvature strongly curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong

Culm - neck medium medium Ear

Plant growth characters

Coleoptile - anthocyanin colouration weak Plant - growth habit intermediate Time of ear emergence late to very late



BROMPTON

CHARACTERS IN DETAIL

Ear

Colour white Length medium lax to medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

short to medium Beak length Beak curvature slightly curved narrow to medium Shoulder width

elevated Shoulder shape

Lower lemma

Beak curvature moderately curved

Grain

red Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence

Flag leaf - anthocyanin of auricles absent or very weak



BUTLER

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile tapering

Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length short

Beak curvature slightly curved Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath absent or very weak
Flag leaf- blade (lower side) absent or very weak
Culm - neck absent or very weak
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



BYSTRA

CHARACTERS IN DETAIL

Ear

Colour: white

Length:short to mediumDensity:medium to denseShape (side profile)parallel sidedAwns or scurs - presence:scurs present

Awns or scurs - distribution:

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness: strong

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak - group 1
Beak length: very short

Beak shape (curvature): slightly curved

Shoulder narrow

Shoulder slightly sloping

Lower Lemma

Beak shape: slightly curved

Grain

Colour red

Colouration with phenol: light to medium

Straw

Length: short Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong
Culm neck: strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin weak
Plant - growth habit: semi-erect
Time ear emergence: medium to late
Flag leaf - Anthocyanin colouration of auricles: absent or very weak



CAMBRIDGE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 3/4
Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium



CASSIUS

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper 1/4

Awns or scurs - length: long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colourationmediumPlant - growth habitintermediateTime of ear emergencelate to very late



CATALYST

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence lat

Flag leaf - anthocyanin of auricles absent or very weak



Variety Name: CELLULE

Crop: Alternative Wheat

Breeder's Reference: fd08114/Cellule **AFP No:**

Breeder: Florimond Desprez Veuve et Fils **Translated from:**

Origin: 15 May 2018

Code: CELL

Coleoptile-anthocyanin colouration: Very strong

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Early

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low

*Flag leaf - Glaucosity of sheath: Very strong

Flag leaf - Glaucosity of leaf blade: Strong ot very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Very thin

*Ear - Glaucosity: Strong to very strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Parallel sided
Ear - Length: Medium to long

*Ear - Density: Dense

*Awns or scurs - presence:

Awns or scurs - distibution:Awns present*Scurs at tip of ear - length:Very long

*Awns at tip of ear - length:

Lower glume - Beak length:Short to medium **Lower glume - Beak shape (curvature):**Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width:Narrow ot medium **Lower glume - Shoulder type:**Slightly sloping or straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs:Weak to medium

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:*Seasonal type:

Alternative

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety.

CHAMPION

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf - blade (lower side) strong to very strong
Culm - neck strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



CHEVRON

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthshort to medium

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence early to medium



CHILTON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributiontip onlyAwns or scurs - length:longApical rachis segment - convex face hairinessmedium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature moderately curved
Shoulder width medium

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium



CHRONICLE

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature slightly curved
Shoulder width very narrow to narrow
Shoulder shape strongly elevated with 2nd pt.

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium



CLAIRE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: short

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



COCOON

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs strong - Group 3

Beak length medium

Beak curvature moderately curved
Shoulder width narrow

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) strong to very strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



COMOROS CS

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



CONQUEROR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



CONSORT

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



CORDIALE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length short

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit semi-erect

Time of ear emergence very early to early
Flag leaf - anthocyanin of auricles absent or very weak



CORONATION

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: long

Awns or scurs - length: long
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong - Group 2+

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence medium to late



COSTELLO

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthshort to medium

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium (1+)

Beak length short

Beak curvature strongly curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



COUGAR

CHARACTERS IN DETAIL

Ear

Colour white
Length short
Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½

Awns or scurs - length: long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



CREEK

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthmedium to longApical rachis segment - convex face hairinessweak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length short

Beak curvature slightly curved Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colourationabsent to weakPlant - growth habitsemi-erectTime of ear emergencemedium to late



CRUSOE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width medium to broad

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath absent or very weak
Flag leaf- blade (lower side) absent or very weak
Culm - neck absent or very weak
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



CUBANITA

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: long

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium to late



DEBEN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



DELPHI

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 3/4
Awns or scurs - length: long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



DENMAN

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: short

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late Flag leaf - anthocyanin of auricles



DEXTER

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium to strong Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium
Beak length medium
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



DICKENS

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong -

Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium



DRUMLIN

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density medium to dense
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length short to medium Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red Colouration with phenol dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



DUNSTON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence very late



DUXFORD

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder shape strongly elevated with 2nd pt.

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) weak
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colourationweakPlant - growth habitsemi-erectTime of ear emergenceearly to medium



EDGAR

scurs present

CHARACTERS IN DETAIL

Ear

Colour:whiteLength:longDensity:laxShape (side profile)tapering

Awns or scurs - presence: Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: weak to medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak - group 1
Beak length: very short to short
Beak shape (curvature): strongly curved

Shoulder narrow

Shoulder straight to elevated

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: medium Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong

Culm neck: strong to very strong
Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin very weak to weak
Plant - growth habit: intermediate
Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



EDMUNDS

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs strong (3)
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



EINSTEIN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length short

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit semi-erect
Time of ear emergence early to medium
Flag leaf - anthocyanin of auricles absent or very weak



ELABORATE

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath weak to medium Flag leaf- blade (lower side) weak to medium

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit intermed. to semi-prostrate

late

Time of ear emergence



ELATION

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to longApical rachis segment - convex face hairinessmedium

Aprical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderately curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

late

Time of ear emergence



ELICIT

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak (group 1)
Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late



ELOQUENCE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium



ELWELL

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium (group 1+)

Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate

Time of ear emergence late



ELYSIUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium
Beak curvature slightly curved
Shoulder width very narrow to narrow
Shoulder shape elev. to strongly elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate

Time of ear emergence late



ENERGISE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence late



EVOLUTION

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½

Awns or scurs - length: long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



EXSEPT

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long lax to medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution tip only Awns or scurs - length: short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved narrow to medium Shoulder width

Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

red Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium Plant - growth habit semi-prostrate Time of ear emergence late to very late Flag leaf - anthocyanin of auricles absent or very weak



FAIRWAY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length very short to short
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface

Extent of internal hairs weak to medium (group 1+)

Beak length short

Beak curvature slightly curved Shoulder width narrow

Shoulder shape slightly sloping

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-prostrate to prostrate

Time of ear emergence medium to late Flag leaf - anthocyanin of auricles



FITZROY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium to dense

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/4

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium (group 1+)

Beak length medium
Beak curvature strongly curved
Shoulder width medium to broad
Shoulder shape straight to elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



FLORETT

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium - group 1 to 2

Beak length: very short to short
Beak shape (curvature): moderately curved

Shoulder narrow

Shoulder slightly sloping to straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: short
Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong

Flag leaf - blade (lower side): weak to medium

Culm neck: medium to strong

Ear - Glaucosity: medium

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: intermediate Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



FOREST

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



FORZOR

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthshort

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak to medium
Culm - neck weak to medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles



FREISTON

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Flag leaf - anthocyanin of auricles

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence very late



Variety Name: Frontal

Crop:

Breeder's Reference: AFP No:

Breeder: Translated from: Dutch

Origin: 24 March 2015

Code: FRNT

Coleoptile-anthocyanin colouration: Absent or very weak

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath:

Low to medium

Strong to very strong

Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Medium *Straw - wall thickness @ 20cms below ear: Very thin

*Ear - Glaucosity: Strong to very strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Tapering

Ear - Length: Short to medium

*Ear - Density: Medium to dense

*Awns or scurs - presence: Present

Awns or scurs - distibution: *Scurs at tip of ear - length:

*Awns at tip of ear - length:

Lower glume - Beak length:

Medium

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: Medium **Lower glume - Shoulder type:** Elevated

Lower glume - External surface roughness: Lower glume - Extent of internal hairs:

Lower glume - Internal imprint:

Lower lemma - Beak shape: Moderately curved

Lower lemma - Beak length-: Medium

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium to strong

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Medium to dark

*Seasonal type: Winter

Extra characters:

30 March 2015



^{* =} Characteristics which should always be included in the description of the variety.

FURLONG

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile tapering scurs present Awns or scurs - presence Awns or scurs - distribution upper 3/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



GALLANT

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium to dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short - medium
Beak curvature slightly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature straight

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitsemi-erectTime of ear emergencemedium



GARIBALDI

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature strongly curved to geniculate

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



GEFION

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long Apical rachis segment - convex face hairiness large Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



Variety Name: Genius

Breeder's Reference: AFP No:

Breeder: Nordsaat Saatzuchtgesellschaft Translated from: German

Crop: Winter Wheat

Origin: Date: 08 October 2010

Code: GENI

Coleoptile-anthocyanin colouration: absent or very weak

*Plant - growth habit: intermediate *Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: very low to low

*Flag leaf - Glaucosity of sheath: strong

Flag leaf - Glaucosity of leaf blade: medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: medium to strong

Hairiness of uppermost node:

*Plant - height: short to medium
*Straw - wall thickness @ 20cms below ear: very thin

*Ear - Glaucosity: weak to medium

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: white

*Ear - Shape (side profile): tapering

Ear - Length: medium to long

*Ear - Density: lax to medium

*Awns or scurs - presence: scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: short

*Awns at tip of ear - length:

Lower glume - Beak length: short

Lower glume - Beak shape (curvature): moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: narrow

Lower glume - Shoulder type: straight to elevated

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: weak - group 1

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: strong *Grain - colour: red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: medium to dark

*Seasonal type: winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety.

GLADIATOR

CHARACTERS IN DETAIL

Ear

Colour white Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence early

Flag leaf - anthocyanin of auricles absent or very weak



GLASGOW

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthshort to mediumApical rachis segment - convex face hairinessweak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence very early to early
Flag leaf - anthocyanin of auricles absent or very weak



GLEAM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length short to medium
Beak curvature v.slightly curved
Shoulder width very narrow to narrow
Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium



GOLDENGUN

CHARACTERS IN DETAIL

Ear

Colour white Length medium

medium to dense Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3 Beak length medium Beak curvature slightly curved Shoulder width narrow

Shoulder shape elev. to elev with 2nd point

Lower lemma

Beak curvature moderately curved

Grain

red Colouration with phenol dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong to very strong Plant - growth habit intermed. to semi-prostrate

late

Time of ear emergence



GRAFTON

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitintermediateTime of ear emergencemedium



Variety Name: Graham Crop:

Breeder's Reference: AFP No:

Breeder: Syngenta Seeds SAS Translated from: French

Origin: Date: 03 November 2014

Code: GRAH

Coleoptile-anthocyanin colouration: Absent or very weak

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath:

Flag leaf - Glaucosity of leaf blade:

Anthers - anthocyanin colouration:

Strong to very strong

Glaucosity of neck:

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Very thin

*Ear - Glaucosity: Strong to very strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile):

Ear - Length:

*Ear - Density:

Parallel sided

Long to very long

Medium to dense

*Awns or scurs - presence: Present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short

*Awns at tip of ear - length:

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature): Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow

Lower glume - Shoulder type: Straight or elevated

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint:

Lower lemma - Beak shape: Slightly curved

Lower lemma - Beak length-: Short

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

Medium

Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 31 March 2015

GRAVITAS

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4

Awns of scurs - distribution upper ¹/₄
Awns or scurs - length: short

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late Flag leaf - anthocyanin of auricles



HAMPDEN

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 3/4 Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium medium to long Beak length Beak curvature moderately curved Shoulder width medium

Shoulder shape elevated

Lower lemma

moderately curved Beak curvature

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak Plant - growth habit intermediate

Time of ear emergence late



HARDWICKE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) weak to medium Culm - neck weak to medium Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence medium to late



HAVANA

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colourationmediumPlant - growth habitsemi-erectTime of ear emergencemedium



HEREFORD

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or Scurs - length: long to very long

Apical rachis segment - convex face hairiness: strong

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length: medium

Beak shape (curvature): strongly curved

Shoulder medium
Shoulder elevated

Lower Lemma

Beak shape: strongly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong

Flag leaf - blade (lower side): strong to very strong

Culm neck: strong
Ear - Glaucosity: strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: intermediate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



HEREWARD

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns of scurs - distributionupper \(\frac{1}{4} \)

Awns or scurs - length: very short to short

Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature slightly curved
Shoulder width broad to very broad

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



HEROLDO

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3
Beak length: short to medium

Beak shape (curvature): straight
Shoulder narrow

Shoulder straight to elevated

Lower Lemma

Beak shape:

Grain

Colour white

Colouration with phenol: light to medium

Straw

Length: short to medium Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): strong to very strong

Culm neck: strong to very strong
Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: semi-prostrate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



HORATIO

CHARACTERS IN DETAIL

Ear

Colour white Length medium lax to medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



HUMBER

absent

CHARACTERS IN DETAIL

Anthers - anthocyanin colouration

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness medium

Lower glume

External surface

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width medium to broad
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate

Time of ear emergence early to medium



Variety Name: **Hybery** Crop: Winter Wheat

Breeder's Reference: surh.3628-164 **AFP No:**

Breeder: Syngenta seeds Ltd **Translated from:**

Origin: 15 May 2018

Code: HYBE

Coleoptile-anthocyanin colouration: Weak

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

Low to medium

*Flag leaf - Glaucosity of sheath:

Flag leaf - Glaucosity of leaf blade:

Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Medium to strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Very thyin

*Ear - Glaucosity: Medium to strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Parallel sided
Ear - Length: Long

*Ear - Density: Medium to dense *Awns or scurs - presence: Scurs present

Awns or scurs - distibution: *Scurs at tip of ear - length: *Awns at tip of ear - length:

Lower glume - Beak length:Very short to short

Lower glume - Beak shape (curvature): Straight to slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width:Narrow **Lower glume - Shoulder type:**Straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak to medium

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:Very weak
*Grain - colour:
Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

Very dark
Winter

Extra characters:

16 May 2018

^{* =} Characteristics which should always be included in the description of the variety.

HYPERION

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthshort

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough
Extent of internal hairs medium (2)
Beak length short to medium
Beak curvature slightly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) Culm - neck

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

strong

Time of ear emergence early to medium
Flag leaf - anthocyanin of auricles absent or very weak



Variety Name: **HYTECK** Crop: Alternative Wheat

Breeder's Reference: hyteck AFP No:

Breeder: Saaten Union Recherche **Translated from:**

Origin: 15 May 2018

Code: HYTE

Coleoptile-anthocyanin colouration: Very weak to weak

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Early to medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low

*Flag leaf - Glaucosity of sheath: Strong to very strong
Flag leaf - Glaucosity of leaf blade: Medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Medium to strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Thin to medium

*Ear - Glaucosity: Medium to strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Parallel sided
Ear - Length: Long to very long

*Ear - Density: Dense

*Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length: Very short to short

Lower glume - Beak shape (curvature): Straight to slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow

Lower glume - Shoulder type: Sloping or slightly sloping

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:Very weak
*Grain - colour:
Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

Dark to very dark
Alternative

Extra characters:

16 May 2018

^{* =} Characteristics which should always be included in the description of the variety.

ICEBREAKER

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium
Beak curvature slightly curved
Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late



ICON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs present

Awns of scurs - distribution upper 3/4
Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium to long
Beak curvature moderately curved
Shoulder width medium

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong

Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



INVICTA

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributiontip onlyAwns or scurs - length:short to media

Awns or scurs - length: short to medium
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late



ISTABRAQ

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper ½Awns or scurs - length:medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect

Time of ear emergence very early to early

Flag leaf - anthocyanin of auricles absent or very weak



JB DIEGO

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution tip only

Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length short

Beak curvature slight to moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

red Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong weak to medium

Ear

Plant growth characters

Coleoptile - anthocyanin colouration strong

semi-erect to intermediate Plant - growth habit

Time of ear emergence medium



JELLICOE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



JORVIK

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape straight

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



JOYFUL

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthlongDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KETCHUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KINGDOM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution tip only
Awns or scurs - length: short
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



KINGFISHER

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long Density medium to dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 medium Awns or scurs - length

weak to medium Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long Beak curvature slightly curved Shoulder width medium elevated Shoulder shape

Lower lemma

Beak curvature moderately curved

Grain

red Colouration with phenol dark

Straw

short to medium Length Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong Culm - neck strong to very strong Ear

strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermed. to semi-prostrate

Time of ear emergence

Flag leaf - anthocyanin of auricles absent or very weak



KIPLING

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitydense to very denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper ½

Awns or scurs - length:

Awns or scurs - length:

Apical rachis segment - convex face hairiness

weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side)

Culm - neck medium
Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KRIS

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium

Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3
Beak length: short to medium
Beak shape (curvature): slightly curved

Shoulder medium

Shoulder straight to elevated

Lower Lemma

Beak shape: slightly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: short
Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): medium to strong

Culm neck: strong
Ear - Glaucosity: strong

Plant growth characters

Coleoptile-anthocyanin medium to strong

Plant - growth habit: semi-prostrate to prostrate

Time ear emergence: late

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



KWS ALABASTER

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length short

Beak curvature slightly curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour white

Colouration with phenol medium to dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence early to medium



KWS BARREL

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence late to very late



KWS BASSET

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length long

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) weak to medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence late to very late



KWS BIELSUM

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitydense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium
A pical rechis segment segment segment medium

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KWS BONHAM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper 1/4Awns or scurs - length:medium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) weak to medium
Culm - neck medium

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late



KWS BRIUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long

Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS BURGHLEY

weak to medium

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Anthers - anthocyanin colouration

Apical rachis segment - convex face hairiness

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

......

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS CASHEL

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length very long
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong

Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath weak

Flag leaf- blade (lower side) very weak to weak

Culm - neck weak Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence



KWS CHILOE

CHARACTERS IN DETAIL

Ear

Colour white Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence awns present Awns of scurs - distribution whole length Awns or scurs - length: very long Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length long to very long
Beak curvature slightly curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol very light to light

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



KWS CLEVELAND

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns of scurs - distribution upper 1/4

Awns or scurs - length: long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Pack langth medium

Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit semi-erect
Time of ear emergence medium



Variety Name: KWS COLOSSEUM Crop: Wheat

Breeder's KWS W370 **AFP No:**

Breeder: KWS UK LTD Translated from:

Date: 28 June 2021 **Origin:**

Code: KCOL

Coleoptile-anthocyanin colouration: Medium to strong

*Plant - growth habit: Intermediate to semi-prostrate

*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low

*Flag leaf - Glaucosity of sheath: Strong to very strong Flag leaf - Glaucosity of leaf blade: Strong to very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Short to medium *Straw - wall thickness @ 20cms below ear: Absent or very thin Strong to very strong

*Ear - Glaucosity:

Ear - Supernumerary spikelets - frequency: *Ear - Colour: White

*Ear - Shape (side profile): Parallel sided Ear - Length: Medium

*Ear - Density: Medium to dense *Awns or scurs - presence: Scur present

Awns or scurs - distibution:

*Scurs at tip of ear - length:

*Awns at tip of ear - length: Medium to long Lower glume - Beak length: Medium

Lower glume - Beak shape (curvature): Slightly curved to moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Medium to broad

Lower glume - Shoulder type: Slightly sloping to straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Medium

Lower glume - Internal imprint: Slightly curved to moderately curved

Lower lemma - Beak shape: Strongly curved to geniculate

Lower lemma - Beak length-:

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Dark to very dark

*Seasonal type: Winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety.

WINTER WHEAT

AFP 1/2416

KWS CONROS

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium (group 1+)

Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong

Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence very late



KWS COYNE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence late

Time of ear emergence Flag leaf - anthocyanin of auricles



KWS CRANIUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence la

Flag leaf - anthocyanin of auricles absent or very weak



KWS CRISPIN

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



KWS CROFT

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: long
Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

•

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



KWS CROSBY

weak

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: long Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2 medium to long Beak length Beak curvature moderately curved narrow to medium Shoulder width

Shoulder shape elevated

Lower lemma

slightly to moderately curved Beak curvature

Grain

Colouration with phenol medium to dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong

Culm - neck strong to very strong Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KWS DALI

CHARACTERS IN DETAIL

Ear

Colour white Length medium to long

Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: medium

weak to medium Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved narrow to medium Shoulder width

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol very dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) weak to medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak Plant - growth habit intermediate Time of ear emergence medium to late



KWS DAWSUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS DELUXUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS DINKUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence lat

Flag leaf - anthocyanin of auricles absent or very weak



KWS EVOKE

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness absent to weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium



KWS EVOLVE

CHARACTERS IN DETAIL

Ear

white Colour Length long Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium
Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence late



KWS EXTASE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlength

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length short

Beak curvature slightly curved Shoulder width slightly curved narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate Time of ear emergence medium



KWS FENWAY

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature INVALID CHARACTER STATE!

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate

Time of ear emergence very late Flag leaf - anthocyanin of auricles



KWS FIREFLY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence late



KWS FOXTUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs weak (group 1)
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium Shoulder shape slightly elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS FULCRUM

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 medium Awns or scurs - length Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS GATOR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS GIGANTUM

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long Density medium to dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 medium Awns or scurs - length Apical rachis segment - convex face hairiness large Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS GUIUM

CHARACTERS IN DETAIL

Ear

white Colour Length long Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4 medium to long Awns or scurs - length weak to medium Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium to strong (group 2+)

Beak length short to medium
Beak curvature strongly curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS HARLYN

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface rough
Extent of internal hairs medium (2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



KWS HENUM

CHARACTERS IN DETAIL

Ear

white Colour Length medium

medium to dense Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long

Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature strongly curved Shoulder width medium to broad

Shoulder shape elev. to strongly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

red Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong Culm - neck strong to very strong Ear

medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence late to very late Flag leaf - anthocyanin of auricles absent or very weak



KWS HORIZON

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness absent to weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitsemi-erectTime of ear emergencemedium to late



KWS JACKAL

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile tapering scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/2 long to very long Awns or scurs - length Apical rachis segment - convex face hairiness weak to medium absent

Anthers - anthocyanin colouration

Lower glume

External surface smooth

medium (group 2) Extent of internal hairs

Beak length medium

moderately curved Beak curvature Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

medium to dark Colouration with phenol

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

absent to weak Coleoptile - anthocyanin colouration Plant - growth habit intermediate medium to late Time of ear emergence



KWS JUBILATION

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length short to medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS JUGUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS KENTON

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium
Culm - neck weak to medium

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS KERANKUM

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) strong
Culm - neck very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS KERRIN

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) weak to medium
Culm - neck weak to medium
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence very late



KWS KIELDER

CHARACTERS IN DETAIL

Ear

Colour white Length medium to long

medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium - Group 2 medium to long Beak length Beak curvature slightly curved very narrow to narrow Shoulder width

elevated Shoulder shape

Lower lemma

slightly curved Beak curvature

Grain

Colouration with phenol

very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit semi-prostrate Time of ear emergence medium to late



KWS KINETIC

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late



KWS KINNELLUM

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 medium to long Awns or scurs - length Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence medium to late Flag leaf - anthocyanin of auricles absent or very weak



KWS LEON

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium (1+)
Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence late to very late



KWS LILI

elevated

CHARACTERS IN DETAIL

Ear

white Colour

Length short to medium

Density dense

Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak absent

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium

Beak length short to medium Beak curvature slightly curved Shoulder width narrow

Shoulder shape

Lower lemma

moderately curved Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



KWS LUMOS

CHARACTERS IN DETAIL

Ear

Colour white

Length
Density
Shape-side profile
Awns or scurs - presence
Awns of scurs - distribution
Awns or scurs - length:
Apical rachis segment - convex face hairiness

short to medium
parallel sided
scurs present
upper \(^1\)4
medium
medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium
Beak curvature strongly curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitsemi-erectTime of ear emergencemedium



KWS LUTHER

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium



KWS MADRYN

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



KWS MCGUGUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length short to medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf - blade (lower side) very strong
Culm - neck very strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KWS MOJITO

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long Density lax to medium parallel sided Shape-side profile Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4

Awns or scurs - length long to very long weak to medium Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium Flag leaf- blade (lower side) weak to medium Culm - neck medium medium Ear

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate late

Time of ear emergence



Variety Name: KWS Montana Crop: Winter wheat

Breeder's Reference: KW 8027-3-08 **AFP No:**

Breeder: KWS LOCHOW GMBH Translated from: German

Date: 22 March 2016 Origin:

Code: KWMON

Coleoptile-anthocyanin colouration: Medium *Plant - growth habit: Semi prostrate

*Time ear emergence: Late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Absent or very low

*Flag leaf - Glaucosity of sheath: Strong Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Short to medium *Straw - wall thickness @ 20cms below ear: Absent or very thin

*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Parallel sided Ear - Length: Medium *Ear - Density: Medium *Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short

*Awns at tip of ear - length:

Lower glume - Beak length: Very short to short

Lower glume - Beak shape (curvature): Straight to slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width:

Lower glume - Shoulder type: Slightly sloping to straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Very weak to weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Strong to very strong

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Medium to dark

*Seasonal type: Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety.





KWS MOZART

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/2 Awns or scurs - length short to medium Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth Extent of internal hairs medium Beak length medium

moderate to strongly curved Beak curvature

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) medium

Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



KWS PALLADIUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlongApical rachis segment - convex face hairinessmedium

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderately curved Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS PARKIN

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthshortDensitydenseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration

Lower glume

External surface entire external surface rough
Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderately curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS PLECTRUM

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature v.slightly curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS PODIUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit semi-erect
Time of ear emergence medium



KWS RACKUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS ROADUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape horizontal to slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS ROWAN

CHARACTERS IN DETAIL

Ear

Colour white Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium



KWS SABRUM

CHARACTERS IN DETAIL

Ear

white Colour Length short Density dense

Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 medium Awns or scurs - length weak to medium Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs strong (group 3) Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

red Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck very strong Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium Plant - growth habit intermediate Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



KWS SANTIAGO

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium - Group 2 Beak length medium

Beak curvature moderately curved narrow to medium Shoulder width

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

red Colouration with phenol medium

Straw

very short to short Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KWS SAXTEAD

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density Shape-side profile tapering Awns or scurs - presence awns present Awns of scurs - distribution whole length Awns or scurs - length: long Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong - Group 3

Beak length long

Beak curvature strongly curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitsemi-erectTime of ear emergencemedium



KWS SCIMITAR

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width broad

Shoulder shape straight to elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence late



KWS SILVERSTONE

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



KWS SILVUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS SISKIN

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium (1+) Beak length short to medium

Beak curvature slight to moderately curved Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong Ear

strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



KWS SKOMER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium to long
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence early



KWS SOLO

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long medium to dense Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4

Awns or scurs - length: medium to long medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium to late



KWS STERLING

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 1/4
Awns or scurs - length: long
Apical rachis segment - convex face hairiness weak

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



KWS TARGET

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ½
Awns or scurs - length: medium
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitsemi-erectTime of ear emergencemedium



KWS TEMPO

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium
Beak length short to medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence very late



KWS TRINITY

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs weak to medium Beak length medium Beak curvature strongly curved Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong

Ear strong

Plant growth characters

Flag leaf - anthocyanin of auricles

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate late

Time of ear emergence



KWS TROMMEL

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature strongly curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence very late



KWS ULTIMATUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)
Beak length medium
Beak curvature strongly curved

Shoulder width very narrow to narrow Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit medium to strong
semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS WEBBUM

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong to very long

Aprical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long
Beak curvature strongly curved
Shoulder width narrow to medium
Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS WRENUM

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitydense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness large to very large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KWS YARIS

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium

Apical rachis segment - convex face hairiness: medium to strong

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length: long

Beak shape (curvature): strongly curved

Shoulder medium
Shoulder elevated

Lower Lemma

Beak shape: strongly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: medium Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong to very strong

Flag leaf - blade (lower side): very strong

Culm neck: strong to very strong
Ear - Glaucosity: strong to very strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: semi-prostrate

Time ear emergence: late

Flag leaf - Anthocyanin colouration of auricles:



KWS ZENTARR

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence late to very late



KWS ZYATT

CHARACTERS IN DETAIL

Ear

white Colour

short to medium Length Density lax to medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs strong (group 3) Beak length short to medium Beak curvature slightly curved Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

red Colour Colouration with phenol dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



LANCASTER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ½Awns or scurs - length:long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck weak to medium
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late



LANGDALE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough
Extent of internal hairs medium (2)
Beak length short to medium
Beak curvature slightly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-erect
Time of ear emergence late



LEAR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



LEEDS

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper ½Awns or scurs - length:long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium



LG ALDANITI

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length very long

Apical rachis segment - convex face hairiness very small to small

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length long to very long

Beak curvature moderate to strongly curved Shoulder width very narrow to narrow

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature

Grain

Colour reddish
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear thick or filled

Glaucosity

Flag leaf - sheath absent or very weak
Flag leaf- blade (lower side) absent or very weak
Culm - neck absent or very weak
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles very weak



LG ASTRONOMER

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long lax to medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4

Awns or scurs - length long to very long medium to strong Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium (group 2) Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong Culm - neck strong to very strong Ear

medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence

Flag leaf - anthocyanin of auricles absent or very weak



LG BLETCHLEY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence late



LG CASSIDY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath absent or very weak
Flag leaf - blade (lower side) absent or very weak
Culm - neck absent or very weak
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



LG DETROIT

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late



LG FARRIER

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface entire external surface rough
Extent of internal hairs medium to strong (group 2+)

Beak length medium
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) strong
Culm - neck very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



LG GENERATION

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)

Beak length medium
Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium Flag leaf- blade (lower side) weak

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence medium to late



LG GILMORE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution tip only

Awns or scurs - length very short to short
Apical rachis segment - convex face hairiness large to very large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length very short to short
Beak curvature slightly curved
Shoulder width narrow to medium
Shoulder shape slightly elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour reddish

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate



LG GRADUATE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length lor

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late



LG ILLUMINATE

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)

Beak length long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence lat

Flag leaf - anthocyanin of auricles absent or very weak



LG INTERSTELLAR

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 medium to long Awns or scurs - length Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs medium (group 2)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate

Time of ear emergence late



LG JIGSAW

CHARACTERS IN DETAIL

Anthers - anthocyanin colouration

Ear

Colour white Length medium Density medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 3/4 Awns or scurs - length long Apical rachis segment - convex face hairiness medium

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)
Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence late to very late



LG MEDUSA

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature strongly curved

Shoulder width very narrow to narrow
Shoulder shape elev. to strongly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



LG MOTOWN

CHARACTERS IN DETAIL

Ear

white Colour Length short Density medium Shape-side profile tapering scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)
Beak length short to medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium

Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



LG NASHVILLE

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 Awns or scurs - length short to medium Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath absent or very weak
Flag leaf- blade (lower side) absent or very weak
Culm - neck absent or very weak
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



LG PHAROS

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long
Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf - blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



LG PRINCE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs weak (group 1)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



LG REDWALD

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitydense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness large
Anthers - anthocyanin colouration absent

Lower glume

External surface

Extent of internal hairs strong (group 3)
Beak length medium to long
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol dark to very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



LG RHYTHM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs weak (group 1)
Beak length medium

Beak curvature moderately curved Shoulder width medium

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late



LG SEEKER

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs medium to strong (group 2+)

Beak length medium
Beak curvature strongly curved
Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit very weak to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



LG SUNDANCE

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthshortDensitydense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) weak
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence very late



LG SABERTOOTH

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium to long Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



LG SKYSCRAPER

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence medium to late



LG SPOTLIGHT

medium to strong

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Anthers - anthocyanin colouration

Apical rachis segment - convex face hairiness

Lower glume

External surface slightly rough
Extent of internal hairs weak (group 1)
Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence late



LG TAPESTRY

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence lat

Flag leaf - anthocyanin of auricles absent or very weak



LG TICKET

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs whole internal surface covered

Beak length medium to long
Beak curvature strongly curved
Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



LG TYPHOON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



LG QUASAR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



LG VOODOO

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath absent or very weak
Flag leaf- blade (lower side) absent or very weak
Culm - neck absent or very weak
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



Lion

CHARACTERS IN DETAIL

Apical rachis segment - convex face hairiness

Ear

white Colour

Length medium to long Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 short to medium Awns or scurs - length small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface

Extent of internal hairs medium to strong (group 2+)

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape horizontal to slightly elevated

Lower lemma

Beak curvature slightly curved

Grain

reddish

Colouration with phenol dark to very dark

Straw

Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong Plant - growth habit intermediate Time of ear emergence early to medium



WINTER WHEAT

AFP 1/2893

LOXTON

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



LUSTRE

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)

Beak length short

Beak curvature slightly curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour white

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence medium



MAGISTER

CHARACTERS IN DETAIL

Ear

white Colour: Length: medium lax to medium Density: Shape (side profile) tapering Awns or scurs - presence: scurs present

Awns or scurs - distribution:

short to medium Awns or scurs - length:

medium Apical rachis segment - convex face hairiness:

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium -group 1 to 2

Beak length: short

Beak shape (curvature): moderately curved Shoulder medium to broad

Shoulder straight

Lower Lemma

Beak shape:

Grain

Colour red Colouration with phenol: dark

Straw

medium to long Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong Flag leaf - blade (lower side): strong Culm neck: strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin weak

semi-prostrate Plant - growth habit: Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



MARIBOSS

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthshortDensitymediumShape-side profiletaperingAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface
Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) strong to very strong

Culm - neck strong

Ear very weak to weak

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence early



MARIS HUNTSMAN

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitydense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium (group 1+)

Beak length medium to long
Beak curvature slightly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length long to very long Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium
Flag leaf - anthocyanin of auricles weak



MARIS WIDGEON

CHARACTERS IN DETAIL

Ear

Colour white Length medium very lax to lax Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs weak - Group 1
Beak length medium
Beak curvature v.slightly curved

Shoulder width v.slightly curved narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very long Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence late
Flag leaf - anthocyanin of auricles weak



Variety Name: Marlowe

Crop:

Breeder's Reference: Sj9726005 **AFP No:** 1/2413

Breeder: Senova Ltd Translated from: Danish

Origin: Date: 23 March 2017

Code: MARL

Coleoptile-anthocyanin colouration:Weak to medium
*Plant - growth habit:
Semi prostrate

*Time ear emergence: Late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Medium to high

*Flag leaf - Glaucosity of sheath: Strong
Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong

Hairiness of uppermost node:

*Plant - height: Medium

*Straw - wall thickness @ 20cms below ear: Absent or very thin

*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White *Ear - Shape (side profile): Parallel

Ear - Length:

*Ear - Density:

*Awns or scurs - presence:

Short to medium

Dense to very dense

Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length:

*Awns at tip of ear - length:

Lower glume - Beak length:

Medium

Long

Lower glume - Beak shape (curvature): Strongly curved

Lower glume - Beak tip:LongLower glume - Shoulder width:MediumLower glume - Shoulder type:Elevated

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Strong

Lower glume - Internal imprint:

Lower lemma - Beak shape:Moderately curved to strongly curved

Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Light to medium

*Seasonal type: Winter

Extra characters:

28 March 2017



^{* =} Characteristics which should always be included in the description of the variety.

MARSTON

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature strongly curved to geniculate

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



Variety Name: Maurizio Crop: Winter Wheat

Breeder's Reference: SZD 6374 **AFP No:**

Breeder: Saatzucht Donau GmbH Translated from: Hungarian

Origin: Date: 19 November 2018

Code: IZIO

Coleoptile-anthocyanin colouration: Abesint or very weak

*Plant - growth habit: Intermediate

*Time ear emergence: Early to medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Very low to low.

*Flag leaf - Glaucosity of sheath: Strong

Flag leaf - Glaucosity of leaf blade: Medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Very strong to strong

Hairiness of uppermost node:

*Plant - height:

*Straw - wall thickness @ 20cms below ear: Thin *Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White *Ear - Shape (side profile): Parallel

Ear - Length:

*Ear - Density:

Lax to medium

*Awns or scurs - presence:

Awns present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Long

*Awns at tip of ear - length:

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature): Moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Medium

Lower glume - Shoulder type: Sloping to sightly sloping

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Medium to dark

*Seasonal type: Winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 19 March 2019

MAXWELL

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presenceawns presentAwns or scurs - distributionwhole lengthAwns or scurs - lengthvery longApical rachis segment - convex face hairinessweak

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length very long

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape strongly elevated with 2nd pt. present

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak to medium
Culm - neck medium

Ear very weak to weak

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence very early



MAYFLOWER

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface
Beak length short to medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) weak to medium

Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong to very strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



MERIT

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long medium to dense Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2

Awns or scurs - length long to very long weak to medium Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium (group 1+)

Beak length

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

red Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong Ear

strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late Flag leaf - anthocyanin of auricles absent or very weak



MINDFUL

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium (group 1+)

Beak length short

Beak curvature strongly curved Shoulder width medium

Shoulder shape horizontal to slightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



WINTER WHEAT

AFP 1/2464

MONROE

CHARACTERS IN DETAIL

Ear

Colour white

Length Short to medium
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

•

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slightly curved

Shoulder width megroup

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate

late

Time of ear emergence



MONTEREY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



Variety Name: MORTIMER Crop: Winter Wheat

Breeder's Reference: SC 2405 **AFP No:**

Breeder: Secobra Translated from:

Origin: 15 May 2018

Code: MORT

Coleoptile-anthocyanin colouration:Absent or very weak*Plant - growth habit:Semi prostrate*Time ear emergence:Medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Very low ot low

*Flag leaf - Glaucosity of sheath: Strong

Flag leaf - Glaucosity of leaf blade: Strong to very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong

Hairiness of uppermost node:

*Plant - height: Short

*Straw - wall thickness @ 20cms below ear: Absent or very thin *Ear - Glaucosity: Medium to strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White
*Ear - Shape (side profile): Parallel
Ear - Length: Long

*Ear - Density: Lax to medium

*Awns or scurs - presence: Present

Awns or scurs - distibution: *Scurs at tip of ear - length:

*Awns at tip of ear - length: Short to medium

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature): Straight to slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow to medium

Lower glume - Shoulder type:

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Very weak to weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak to medium

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:*Seasonal type:

Medium
Winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety.

MOSAIC

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering scurs present Awns or scurs - presence Awns or scurs - distribution upper 3/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit medium to strong semi-erect to intermediate

Time of ear emergence late to very late



Variety Name: Moschus Crop: Winter whrat

Breeder's Reference: STRU 114017 **AFP No:**

Breeder: Dr HermannStrube Translated from: German

Origin: Date: 19 November 2018

Code: MOSC

Coleoptile-anthocyanin colouration: Strong to very strong

*Plant - growth habit: Intermediate
*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath:

Flag leaf - Glaucosity of leaf blade:

Strong to very strong

Strong to very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Medium

*Straw - wall thickness @ 20cms below ear: Absent or very thin

*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour:

*Ear - Shape (side profile):

Ear - Length:

*Ear - Density:

*Awns or scurs - presence:

Tapering

Medium to long

Lax to medium

Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature): Slightly to moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow to medium

Lower glume - Shoulder type: Straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint: Lower lemma - Beak shape:

Lower lemma - Beak length-:

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium to strong

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:*Seasonal type:

Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 19 March 2019



MOULTON

CHARACTERS IN DETAIL

Ear

white Colour Length long Density medium Shape-side profile tapering scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late



MV FREDERICIA

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presenceawns presentAwns or scurs - distributionwhole lengthAwns or scurs - lengthvery long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length long to very long

Beak curvature moderate to strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



MYRIAD

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¾

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



MYSTIC CS

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium to late



NELSON

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



OAKLEY

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium



OLIVIN

CHARACTERS IN DETAIL

Ear

Colour:whiteLength:mediumDensity:mediumShape (side profile)parallel sidedAwns or scurs - presence:scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: medium to strong

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: medium - group 2
Beak length: very short to short
Beak shape (curvature): strongly curved
Shoulder narrow to medium

Shoulder slightly sloping to straight

Lower Lemma

Beak shape: strongly curved to geniculate

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: medium to long

Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): medium
Culm neck: medium

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin strong to very strong

Plant - growth habit: intermediate to semi-prostrate

Time ear emergence: late

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



OLYMPUS

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4

Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



Variety Name: Opus

Breeder's Reference: PBIS 99/70 AFP No:

Breeder: Erhardt Eger Translated from: German

Crop: Winter Wheat

Date: 23 November 2001 **Origin:**

Code: OPUS

Coleoptile-anthocyanin colouration: absent or very weak

*Plant - growth habit: intermediate to semi-prostrate

*Time ear emergence: medium to late Flag leaf - Anthocyanin colouration of auricles: absent or very weak

Frequency of plants with recurved flag leaves: very low to low

*Flag leaf - Glaucosity of sheath: strong Flag leaf - Glaucosity of leaf blade: medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: strong

Hairiness of uppermost node:

*Plant - height: medium *Straw - wall thickness @ 20cms below ear: very thin

*Ear - Glaucosity: medium

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: white *Ear - Shape (side profile): tapering Ear - Length: medium *Ear - Density: medium

*Awns or scurs - presence: scurs present

Awns or scurs - distibution: *Scurs at tip of ear - length: short *Awns at tip of ear - length:

Lower glume - Beak length: Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: Lower glume - Shoulder type:

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs:

Lower glume - Internal imprint:

Lower lemma - Beak shape: Lower lemma - Beak length-:

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour: **Grain - Shape:**

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type: winter

Extra characters:

* = Characteristics which should always be included in the description of the variety.

06 May 2020



ORATOR

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletapering

Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/4

Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)
Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) weak to medium
Culm - neck medium to strong
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-prostrate

Time of ear emergence medium to late



ORBIT

CHARACTERS IN DETAIL

Ear

white Colour Length long Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short to medium Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



OSMIUM

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthshortDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak (group 1)

Beak length short

Beak curvature v.slightly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



OXFORD

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish Colouration with phenol dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-prostrate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



PANACEA

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¾Awns or scurs - length:long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late



PANORAMA

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length: short to medium
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Ear

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) weak to medium

Culm - neck medium

Plant growth characters
Coleoptile - anthocyanin colouration

Plant - growth habit semi-erect to intermediate

weak to medium

strong

Time of ear emergence late



PHILOSOPHER

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length: long
Apical rachis segment - convex face hairiness weak

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



PIRENEO

weak to medium

CHARACTERS IN DETAIL

Ear

Colour: white Length: medium

Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: awns present

Awns or scurs - distribution:

Awns or scurs - length: long
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: medium - group 2
Beak length: long to very long
Beak shape (curvature): slightly curved

Shoulder narrow Shoulder straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: medium

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Culm neck:

Flag leaf - sheath: weak

Flag leaf - blade (lower side): very weak to weak

Ear - Glaucosity: weak

Plant growth characters

Coleoptile-anthocyanin

Plant - growth habit: intermediate
Time ear emergence: early to late

Flag leaf - Anthocyanin colouration of auricles:



Variety Name: **PONTICUS** Crop: Winter Wheat

Breeder's Reference: STRU 114010 **AFP No:**

Breeder: Dr Hermann Strube **Translated from:** German

Origin: Date: 14 November 2017

Code: PONT

Coleoptile-anthocyanin colouration: Strong

*Plant - growth habit: Intermediate to semi-prostrate

*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath:

Very strong

Very strong

Very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Very strong

Hairiness of uppermost node:

*Plant - height: Strong to medium
*Straw - wall thickness @ 20cms below ear: Absent or very thin

*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Tapering

Ear - Length: Medium to long

*Ear - Density: Lax

*Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length: Very short to short

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width:Narrow to medium **Lower glume - Shoulder type:**Slightly sloping

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

Medium

Winter

Extra characters:

28 March 2018

^{* =} Characteristics which should always be included in the description of the variety.

Variety Name: PRAKTIK

Crop:

Breeder's Reference: AFP No: 1/

Breeder: RAGT Translated from:

Origin: GERMANY Date: 22 October 2012

Code: PRAK

Coleoptile-anthocyanin colouration: ABSENT OR VERY WEAK

*Plant - growth habit: INTERMEDIATE TO SEMI-PROSTRATE

*Time ear emergence: MEDIUM TO LATE

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

ABSENT OR VERY LOW

*Flag leaf - Glaucosity of sheath: MEDIUM
Flag leaf - Glaucosity of leaf blade: MEDIUM

Anthers - anthocyanin colouration:

Glaucosity of neck: MEDIUM TO STRONG

Hairiness of uppermost node:

*Plant - height: SHORT

*Straw - wall thickness @ 20cms below ear: very thin

*Ear - Glaucosity: MEDIUM

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: WHITE *Ear - Shape (side profile): TAPERING

Ear - Length: SHORT TO MEDIUM

*Ear - Density: MEDIUM

*Awns or scurs - presence: SCURS PRESENT

Awns or scurs - distibution:

*Scurs at tip of ear - length: VERY SHORT TO SHORT

*Awns at tip of ear - length:

Lower glume - Beak length: VERY SHORT TO SHORT

Lower glume - Beak shape (curvature): STRAIGHT

Lower glume - Beak tip:SLIGHTLY CURVEDLower glume - Shoulder width:NARROW TO MEDIUM

Lower glume - Shoulder type:

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: MEDIUM

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour:

*RED

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: DARK

*Seasonal type: WINTER TYPE

Extra characters: STRAW:PILTH IN CROSS SECTION -

VERY THIN

30 March 2015



^{* =} Characteristics which should always be included in the description of the variety.

QI

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - length:medium

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit semi-erect
Time of ear emergence medium



RA3W 08

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness: weak to medium

Anthers - anthocyanin

Lower Glume

External surface roughness: medium

Extent of internal hairs: medium - group 2

Beak length: medium

Beak shape (curvature): moderately curved

Shoulder broad
Shoulder elevated

Lower Lemma

Beak shape: moderately curved to strongly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: short to medium

Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): medium to strong

Culm neck: weak
Ear - Glaucosity: medium

Plant growth characters

Coleoptile-anthocyanin medium
Plant - growth habit: intermediate
Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



RAMSAY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)

Beak length short

Beak curvature slightly curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



RAPSODIA

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium

Density: medium
Shape (side profile) tapering

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: weak to medium

Anthers - anthocyanin

Lower Glume

External surface roughness: smooth

Extent of internal hairs: medium to strong - group 2 to 3

Beak length: short

Beak shape (curvature): moderately curved

Shoulder medium Shoulder straight

Lower Lemma

Beak shape: slightly curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: short

Wall thickness @ 20cms below ear: very thin to thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): medium to strong

Culm neck: medium to strong
Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin strong

Plant - growth habit: semi-prostrate to prostrate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



RAZZANO

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs – distribution

Awns or scurs - length: very short
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface
Extent of internal hairs medium to strong - Group 2+

Beak length short

Beak curvature v.slightly curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



REFLECTION

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution upper 3/4
Awns or scurs - length medium
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence medium to late



RELAY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium medium to dense Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2

Awns or scurs - length: long

Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium Beak curvature slightly curved Shoulder width narrow Shoulder shape elevated

Lower lemma

slightly curved Beak curvature

Grain

red Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

strong Ear

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



REVELATION

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



RGT ADVENTURE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length short to medium
Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



RGT ANELLE

CHARACTERS IN DETAIL

Ear

Colour coloured
Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length very short to short
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck strong to very strong
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



RGT BAIRSTOW

CHARACTERS IN DETAIL

Ear

white Colour

short to medium Length Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 medium to long Awns or scurs - length very strong

Apical rachis segment - convex face hairiness absent Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong Culm - neck strong to very strong Ear

medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak intermediate Plant - growth habit Time of ear emergence medium to late Flag leaf - anthocyanin of auricles absent or very weak



RGT BLOSSOM

weak to medium

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 medium to long Awns or scurs - length

Anthers - anthocyanin colouration

Apical rachis segment - convex face hairiness

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length short to medium Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol medium

Straw

short to medium Length Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong medium to strong

Ear

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong Plant - growth habit intermediate Time of ear emergence late



RGT BOWEN

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlongApical rachis segment - convex face hairinessweak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length shor

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate

Time of ear emergence late



RGT COOK

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presenceawns presentAwns or scurs - distributionwhole lengthAwns or scurs - lengthvery long

Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs strong to whole surface

Beak length long to very long
Beak curvature strongly curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf - blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



RGT CONVERSION

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length long

Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence late



RGT DRAX

absent or very weak

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitydenseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Anthers - anthocyanin colouration absent

Apical rachis segment - convex face hairiness

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



RGT FLINTOFF

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presenceawns presentAwns or scurs - distributionwhole lengthAwns or scurs - lengthvery long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs whole internal surface covered

Beak length very long

Beak curvature moderate to strongly curved Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



RGT GALACTUS

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



Variety Name: RGT GRADIENT Crop: WW

Breeder's Reference: RW41394 **AFP No:**

Breeder: R2n sas **Translated from:** Danish

Origin: Trident + Gravitas Date: 22 March 2016

Code: RGTG

Coleoptile-anthocyanin colouration: Very weak to weak

*Plant - growth habit: Prostrate
*Time ear emergence: Medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low to medium

*Flag leaf - Glaucosity of sheath: Strong

Flag leaf - Glaucosity of leaf blade: Medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Absent or Very thin

*Ear - Glaucosity: Medium to strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Tapering

Ear - Length: Short to medium

*Ear - Density: Medium
*Awns or scurs - presence: Present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Long

*Awns at tip of ear - length:

Lower glume - Beak length:Medium to long **Lower glume - Beak shape (curvature):**Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Medium **Lower glume - Shoulder type:** Elevated

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Medium

Lower glume - Internal imprint:

Lower lemma - Beak shape: Moderately curved

Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak to medium

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

Dark

Winter

Extra characters:

28 March 2018

^{* =} Characteristics which should always be included in the description of the variety.

RGT GRAVITY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium Density medium

Shape-side profile tapering scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/2 Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

medium (group 2) Extent of internal hairs

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



RGT HOWLEY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness large to very large

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-prostrate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



RGT ILLUSTRIOUS

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium Apical rachis segment - convex face hairiness strong Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium to strong (2+)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



RGT KNIGHTSBRIDGE

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) weak
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence very late



RGT KOI

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 medium to long Awns or scurs - length

Apical rachis segment - convex face hairiness strong Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium Beak curvature strongly curved

Shoulder width narrow

Shoulder shape slightly sloping

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Glaucosity

short to medium Length Wall thickness @20cms below ear absent or very thin

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong

Culm - neck strong Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late Flag leaf - anthocyanin of auricles absent or very weak



RGT LANTERN

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution
Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



RGT MARLBOROUGH

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)
Beak length medium to long
Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence late to very late



RGT NONU

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to longApical rachis segment - convex face hairinessmedium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs strong to whole surface

Beak length medium
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape horizontal to slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol dark to very dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit semi-prostrate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT PADDINGTON

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved Shoulder width marrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) weak to medium
Culm - neck medium

Ear medium medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



RGT PEMBROKE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate

Time of ear emergence late to very late



RGT QUICKSILVER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT RASHID

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence very late

Flag leaf - anthocyanin of auricles absent or very weak



RGT SACRAMENTO

CHARACTERS IN DETAIL

Ear

white Colour Length short Density medium Shape-side profile tapering awns present Awns or scurs - presence whole length Awns or scurs - distribution Awns or scurs - length very long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length long to very long
Beak curvature slightly curved
Shoulder width very narrow to narrow
Shoulder shape slightly sloping

Lower lemma

Beak curvature slightly curved

Grain

Colour red Colouration with phenol light

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



RGT SAKI

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length long

Beak curvature strongly curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT SILVERSURFER

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature moderately curved Shoulder width narrow to medium Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT STOKES

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT VENOM

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT WASABI

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf - blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT WILKINSON

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 medium Awns or scurs - length Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs medium (group 2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-prostrate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT WOLVERINE

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short strong Apical rachis segment - convex face hairiness Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)

Beak length short

Beak curvature slightly curved Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT ZINZAN

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs strong (group 3)
Beak length short to medium

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



SARTORIAL

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlongApical rachis segment - convex face hairinessstrong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs weak (group 1)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence late

Time of ear emergence Flag leaf - anthocyanin of auricles



SY COACH

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length very short to short
Apical rachis segment - convex face hairiness large to very large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium
Beak curvature strongly curved
Shoulder width medium

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



RGT SCRUMMAGE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium to long
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium
Apical rachis segment - convex face hairiness weak

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence late



RGT SORCERER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profiletaperingAwns or scurs - presenceawns present

Awns or scurs - distribution

Awns or scurs - length n/a

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



RGT WESTMINSTER

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present

Awns or scurs - distribution

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) weak to medium

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence very late



WINTER WHEAT

AFP 1/2519

RGT UNIVERSE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



RIBAND

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs weak (1)
Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) weak
Culm - neck medium
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak

Plant - growth habit prostrate
Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



ROADSTER

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presenceawns present

Awns or scurs - distribution

Awns or scurs - length n/a
Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length long to very long
Beak curvature moderately curved
Shoulder width very narrow to narrow

Shoulder shape straight

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red Colouration with phenol light

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



ROBIGUS

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium
Flag leaf - anthocyanin of auricles weak



ROWLING

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



ROYSTON

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)
Beak length short to medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) weak to medium
Culm - neck weak to medium
Ear medium

Plant growth characters

Flag leaf - anthocyanin of auricles

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence late



RUSKIN

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium
Beak length short to medium
Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



SAHARA

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ½

Awns or scurs - length:

Apical rachis segment - convex face hairiness

weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak

Culm - neck weak

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



SAVELLO

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



SCOUT

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature straight to slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



SCRIBE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium Beak curvature slightly curved Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

red

Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong Culm - neck strong Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate

Time of ear emergence medium



SHABRAS

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium
Beak curvature moderately curved

Shoulder width marrow to medium

Shoulder shape strongly elevated with 2nd pt. present

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



SHEPHERD

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderate to strongly curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late



SHERLOCK

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium
Shore side profile

Shape-side profile tapering
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length very long

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length long to very long

Beak curvature moderate to strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



SHIRE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness medium

Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



SKYFALL

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence awns present Awns or scurs - distribution whole length very long Awns or scurs - length: Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length long to very long
Beak curvature v.slightly curved
Shoulder width very narrow to narrow
Shoulder shape elev. to elev with 2nd point

Lower lemma

Beak curvature straight

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium



SMUGGLER

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration very strong
Plant - growth habit intermediate
Time of ear emergence early

Flag leaf - anthocyanin of auricles absent or very weak



SOISSONS

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length:

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak - Group 1

Beak length

Beak curvature

Shoulder width narrow

Shoulder shape slightly sloping

Lower lemma

Beak curvature straight

Grain

Colour red

Colouration with phenol

Straw

Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath absent or very weak

Flag leaf- blade (lower side) weak

Culm - neck absent or very weak Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence very early to early
Flag leaf - anthocyanin of auricles absent or very weak



SOLACE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ½Awns or scurs - length:long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



SOLSTICE

CHARACTERS IN DETAIL

Ear

Colour white Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 medium to long Awns or scurs - length: Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence late
Flag leaf - anthocyanin of auricles very weak



SOLTEK

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density dense Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short to medium Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium
Beak curvature moderately of

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



Variety Name: **SORRIAL** Crop: WINTER WHEAT

Breeder's Reference: AFP No:

Origin: FRANCE Date: 03 June 2013

Code: SORRI

Coleoptile-anthocyanin colouration: MEDIUM

*Plant - growth habit: INTERMEDIATE TO SEMI-PROSTRATE

*Time ear emergence: EARLY TO MEDIUM

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: VERY LOW TO LOW *Flag leaf - Glaucosity of sheath: VERY STRONG

Flag leaf - Glaucosity of leaf blade: STRONG TO VERY STRONG

Anthers - anthocyanin colouration:

Glaucosity of neck: STRONG TO VERY STRONG

Hairiness of uppermost node:

*Plant - height: SHORT
*Straw - wall thickness @ 20cms below ear: VERY THIN

*Ear - Glaucosity: STRONG TO VERY STRONG

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: WHITE

*Ear - Shape (side profile): PARALLEL SIDED
Ear - Length: MEDIUM TO LONG

*Ear - Density: DENSE

*Awns or scurs - presence: SCURS PRESENT

Awns or scurs - distibution:

*Scurs at tip of ear - length: VERY SHORT TO SHORT

*Awns at tip of ear - length: Lower glume - Beak length:

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: MEDIUM TO BROAD

Lower glume - Shoulder type: STRAIGHT

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: WEAK

Lower glume - Internal imprint:

Lower lemma - Beak shape: SLIGHTLY CURVED

Lower lemma - Beak length-: SHORT

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: WEAK TO MEDIUM

*Grain - colour: RED

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

WEDIUM
WINTER

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015

SPEARHEAD

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface

Beak length short to medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



SPOCK

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late to very late



SPYDER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



STIGG

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensityvery denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length: short to medium
Apical rachis segment - convex face hairiness weak

Apical rachis segment - convex face nairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



STRATOSPHERE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved Shoulder width marrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) weak
Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



SWALLOW

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length long

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence lat

Flag leaf - anthocyanin of auricles absent or very weak



SY EPSON

CHARACTERS IN DETAIL

Ear

Colour: white Length: long

Density: lax to medium
Shape (side profile) parallel sided
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: long
Apical rachis segment - convex face hairiness: very weak

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium - group 1 to 2

Beak length: short to medium

Beak shape (curvature): slightly to moderately curved

Shoulder narrow to medium

Shoulder slightly sloping or straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: medium

Straw

Length:

Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: medium to strong

Flag leaf - blade (lower side): medium

Culm neck: medium to strong
Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin absent to very weak Plant - growth habit: semi-prostrate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



SY GRIFFIN

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence late to very late



SY INSITOR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence la

Flag leaf - anthocyanin of auricles absent or very weak



SY LOKI

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence medium to late



SY MEDEA

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmediumApical rachis segment - convex face hairinessstrong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface
Beak length very short to short
Beak curvature slightly curved
Shoulder width medium to broad

Shoulder shape elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath absent or very weak
Flag leaf- blade (lower side) absent or very weak
Culm - neck absent or very weak
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium



THEODORE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



TIMARU

CHARACTERS IN DETAIL

Ear

Colour white Length long dense Density

Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution tip only Awns or scurs - length: very short Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium Beak curvature strongly curved narrow to medium Shoulder width

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colouration with phenol light to medium

Straw

medium Length Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong Ear medium to strong

Plant growth characters Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



TOBY

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density dense to very dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short
Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermed. to semi-prostrate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



TORCH

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution upper 1/4
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



TORPHINS

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium to late



TREASURE

CHARACTERS IN DETAIL

Ear

Colour white

short to medium Length Density medium to dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long medium Apical rachis segment - convex face hairiness Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



TUXEDO

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



TWISTER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ½

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium



UNIVERSAL

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium to long
Beak curvature strongly curved
Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence late



VELOCITY

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributiontip only

Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness medium

Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature strongly curved to geniculate

Shoulder width medium to broad

Shoulder shape strongly elevated with 2nd pt. present

Lower lemma

Beak curvature geniculate

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) weak to medium

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence early to medium



VERSO

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature v.slightly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature straight to slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



VISCOUNT

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitydenseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - presence scurs present

Awns or scurs - distribution upper \(^{1}\)4

Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



Variety Name: Vulkan Crop: Winter Wheat

Breeder's Reference: AFP No:

Origin: 11 March 2014

Code: VULK

Coleoptile-anthocyanin colouration:

*Plant - growth habit: Semi erect to intermediate

*Time ear emergence: Very early to early

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath:

Low

Medium

Flag leaf - Glaucosity of leaf blade: Very weak to weak

Anthers - anthocyanin colouration:

Glaucosity of neck: Weak to medium

Hairiness of uppermost node:

*Plant - height: Medium

*Straw - wall thickness @ 20cms below ear: Very thick or filled

*Ear - Glaucosity:

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Parallel sided
Ear - Length: Very short

*Ear - Density: Medium

*Awns or scurs - presence: Awns or scurs - distibution:

*Scurs at tip of ear - length:

*Awns at tip of ear - length:

Medium

Medium

Lower glume - Beak length:Lower glume - Beak shape (curvature):
Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Absent or very narrow

Present

Lower glume - Shoulder type: Sloping

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Medium

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type: Winter

Extra characters:



WARRIOR

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear n/a

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



WEAVER

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length: long
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit intermediate
Time of ear emergence medium



WELCOME

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presenceawns presentAwns or scurs - distributionwhole lengthAwns or scurs - lengthvery long

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length very long

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape elev. to elev with 2nd point

Lower lemma

Beak curvature straight

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



WELFORD

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



XI19

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long

Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - length: short

Awns or scurs - distribution tip only
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles very weak



ZEBEDEE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - length:short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



ZOOM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)
Beak length medium to long
Beak curvature strongly curved

Shoulder width narrow

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



ZULU

CHARACTERS IN DETAIL

Ear

Colour white Length medium medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4 Awns or scurs - length: long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red Colouration with phenol dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) weak to medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermed. to semi-prostrate
Time of ear emergence medium to late

