

WINTER BARLEY

AFP 2/2073

ACCRUE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	short to medium
Shape	parallel

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	short
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	medium
Awn tips	medium to strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late to very late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/2586

ACUTE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	medium
Awn tips	weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

AMARENA**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium to dense
Length	medium to long
Shape	parallel

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	short
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

Variety Name: **AMISTAR**

Crop: Winter barley

Breeders Reference: MH 05 AM 04

AFP No:

Breeder: SARL

Translated from: French

Origin:

Date: 27 March 2018

Code: AMIS

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Early
*Flag leaf- anthocyanin colouration of auricles:	Medium to strong
Frequency of plants with recurved flag leaves:	Medium
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Medium
Ear- attitude:	Erect to semi erect
Ear- length:	Medium to long
*Ear- number of rows:	more than two
*Ear- density:	Medium to dense
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Medium
*Awn- length compared to ear:	Medium to long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	Medium
Collar type:	
Rachis- length of first segment:	Short to medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Absent or very weak to weak
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Strong
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Medium to strong
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

28 March 2018

WINTER BARLEY

AFP 2/2447

ARCHER

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax to medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	long
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	semi-prostrate to prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

BAZOOKA**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	short
Shape	parallel

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	short to medium
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	very weak to weak
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

BELFRY**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	short
Shape	parallel

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	short to medium
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2375

BLAZING

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	long
Shape	

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

BLUEBELL**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	medium
Shape	parallel

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium to strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect

WINTER BARLEY

AFP 2/2581

CADILLAC

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	very weak to weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2448

CALIFORNIA

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	long
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/1825

CAMION

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax to medium
Length	long
Shape	tapering

Awn

Length compared to ear	+/- equal to longer
------------------------	---------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	weak
Awn tips	weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	very early to early
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

CAMPANILE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium to dense
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short (shorter than ear)
------------------------	--------------------------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium to strong
Rachilla hair type	short
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	horizontal to semi-recurved

WINTER BARLEY

AFP 2/1606

CARAT

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax to medium
Length	medium
Shape	tapering

Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

Rachis

Length of first segment	short to medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	horizontal to semi-recurved

Variety Name: **CARIBIC**

Crop: Winter Barley

Breeders Reference: NIC 07-5440-C

AFP No:

Breeder: Limagrain GmbH

Translated from: German

Origin:

Date: 27 March 2018

Code: CARB

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Absent
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	Low to medium
Flag leaf- glaucosity of sheath:	Medium
*Ear- glaucosity:	Very weak to weak
Ear- attitude:	Horizontal
Ear- length:	Short to medium
*Ear- number of rows:	Two
*Ear- density:	Lax
Ear- shape:	Tapering
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Short to medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	NA
Sterile spikelet-shape of tip:	NA
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	
Collar type:	
Rachis- length of first segment:	Medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium to strong
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

28 March 2018

WINTER BARLEY

AFP 2/2058

CASSATA

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium to dense
Length	medium
Shape	tapering

Awn

Length compared to ear	+/- equal to longer
------------------------	---------------------

Rachis

Length of first segment	short to medium
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	absent (grains thresh free)
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	medium to strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/2582

CAVALIER

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	medium to strong
Lemma nerves - at early dough	strong to very strong
Awn tips	strong to very strong

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2657

CELOONA

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	dense
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	short to medium
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	medium

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emergence.	semi-erect

Variety Name: **CENTAURE**

Crop: Winter Barley

Breeders Reference: sc30007ph

AFP No:

Breeder: Secobra Recherches

Translated from:

Origin:

Date: 15 May 2018

Code: AURE

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong to very strong
*Ear- glaucosity:	Medium
Ear- attitude:	Semi erect to horizontal
Ear- length:	Medium
*Ear- number of rows:	Non or rudimentary
*Ear- density:	Dense
Ear- shape:	Tapering to parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	NA
Sterile spikelet-shape of tip:	NA
Sterile spikelet- length of lemma:	NA
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	Short to medium
Collar type:	
Rachis- length of first segment:	Short
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Weak to medium
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

16 May 2018

WINTER BARLEY

AFP 2/2887

COREF

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	medium to strong
Ear - attitude	semi-erect

WINTER BARLEY

AFP 2/2743

CRAFT

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	strong
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/1983

CYPRESS

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	medium
Lemma nerves - at early dough	strong
Awn tips	strong

Plant growth characters

Plant - growth habit	semi-prostrate to prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-recurved

Variety Name: **Daxor**

Crop: Winter Barley

Breeders Reference:

Breeder: Sarl Adrien Momont et Fils

Origin:

Code: DAXO

AFP No:

Translated from: French

Date: 24 March 2015

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Absent
*Time of ear emergence:	
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	Low
Flag leaf- glaucosity of sheath:	Medium to strong
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Semi- erect
Ear- length:	Medium
*Ear- number of rows:	More than two
*Ear- density:	Medium to dense
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	
*Awn- length compared to ear:	Medium to long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	Absent to very weak
*Plant height:	
Collar type:	
Rachis- length of first segment:	Short to medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Weak
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Strong
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

31 March 2015

ELECTRUM**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	weak to medium
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude	erect to semi-erect

WINTER BARLEY

AFP 2/2368

ELEMENT

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	medium
Length	medium to long
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	very weak to weak
Lemma nerves - at early dough	weak to medium
Awn tips	very weak to weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

WINTER BARLEY

AFP 2/2383

ESCADRE

CHARACTERS IN DETAIL

Ear

Number of rows	six
Density	dense
Length	short
Shape	parallel

Awn

Length compared to ear	+/- equal to longer
------------------------	---------------------

Rachis

Length of first segment	short to medium
Curvature of first segment	very weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	weak to medium
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

Variety Name: **Estoria**

Crop: Winter Barley

Breeders Reference: SE 213/10 WG

AFP No: 2/

Breeder: Landwirtschaftliche Fachschule
Edelhof - Saatzucht

Translated from: German

Origin: Austria

Date: 12 May 2017

Code: ESTO

*Plant - growth habit:	semi prostrate
*Lower leaves - hairiness of leaf sheaths:	present
*Time of ear emergence:	medium
*Flag leaf- anthocyanin colouration of auricles:	absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	weak to medium
*Ear- glaucosity:	weak
Ear- attitude:	semi erect
Ear- length:	medium to long
*Ear- number of rows:	two
*Ear- density:	dense to very dense
Ear- shape:	tapering
*Awn- anthocyanin colouration of the tips:	absent or very weak
*Awn- length compared to ear:	medium to long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	n/a
Sterile spikelet-shape of tip:	n/a
Sterile spikelet- length of lemma:	n/a
Median spikelet- length of glume + awn cf grain:	equal
Lemma nerves- anthocyanin colouration:	absent or very weak
*Plant height:	medium
Collar type:	
Rachis- length of first segment:	short
Rachis- width of first segment:	
Rachis- curvature of first segment:	weak
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	whitish
*Grain- rachilla hair type:	long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	absent or very weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	
*Grain- ventral furrow - presence of hairs:	absent
*Grain- husk:	present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	collar
*Seasonal type:	winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

28 March 2018

WINTER BARLEY

AFP 2/2895

FANION

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	medium
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	longer

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect

WINTER BARLEY

AFP 2/2596

FENTARA

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium to long
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

Variety Name: **Findora**

Crop: Winter Barley

Breeders Reference:

AFP No: 2/2676

Breeder:

Translated from: German

Origin: Germany

Date: 09 April 2015

Code: FIND

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Early
*Flag leaf- anthocyanin colouration of auricles:	Absent or very week
Frequency of plants with recurved flag leaves:	Low
Flag leaf- glaucosity of sheath:	Medium
*Ear- glaucosity:	Medium
Ear- attitude:	Semi-erect to horizontal
Ear- length:	Medium
*Ear- number of rows:	Two
*Ear- density:	Dense
Ear- shape:	Tapering
*Awn- anthocyanin colouration of the tips:	Absent or very week
*Awn- length compared to ear:	Medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	N/A - non or rudimentary
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	Equal
Lemma nerves- anthocyanin colouration:	Absent or very week
*Plant height:	Very short to short
Collar type:	
Rachis- length of first segment:	Short to medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Short
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Very week to week
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very week
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

09 April 2015

WINTER BARLEY

AFP 2/1910

FLAGON

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax to medium
Length	long
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	horizontal to semi-recurved

WINTER BARLEY

AFP 2/2664

FLETCHER

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	dense
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	short to medium
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2392

FLORENTINE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	medium
Lemma nerves - at early dough	medium
Awn tips	medium

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2807

FUNKY

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	dense
Length	short
Shape	parallel

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	short
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	strong
Ear - attitude 21 days after ear emerg.	semi-erect

GALATION**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	short to medium
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	weak to medium
Awn tips	weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/2583

HARLEQUIN

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	shorter to +/- equal
------------------------	----------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

WINTER BARLEY

AFP 2/2821

JETTOO

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	medium
Length	short to medium
Shape	parallel

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	medium
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	longer

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	medium
Awn tips	medium

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude	semi-erect to horizontal

WINTER BARLEY

AFP 2/2755

KATHMANDU

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short
------------------------	-------

Rachis

Length of first segment	medium to long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early
Flag leaf - glaucosity of sheath	weak
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

Variety Name: **KWS Astaire**

Crop:

Breeders Reference: KW6-341

AFP No: 2/2877

Breeder: KWS UK Ltd

Translated from: German

Origin:

Date: 23 March 2017

Code: KAST

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong to very strong
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Semi erect
Ear- length:	Medium
*Ear- number of rows:	More than two
*Ear- density:	Medium
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Medium to long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Very long
Lemma nerves- anthocyanin colouration:	
*Plant height:	Medium
Collar type:	
Rachis- length of first segment:	Medium to long
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium to strong
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Strongly coloured
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Medium
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

28 March 2017

WINTER BARLEY

AFP 2/2650

KWS CAPELLA

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	very lax to lax
Length	short to medium
Shape	tapering

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	very weak to weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2802

KWS CARBIS

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect

KWS CARIBOU**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax to medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect

WINTER BARLEY

AFP 2/2309

KWS CASSIA

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	short
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	very weak to weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

WINTER BARLEY

AFP 2/2804

KWS CRESWELL

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	medium to long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect

KWS FARO**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium to dense
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	shorter

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium to strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	very weak to weak
Awn tips	weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	weak
Ear - attitude	erect to semi-erect

KWS GIMLET**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	very lax to lax
Length	long to very long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude	horizontal

WINTER BARLEY

AFP 2/2523

KWS GLACIER

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/2650

KWS INFINITY

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	very lax to lax
Length	short to medium
Shape	tapering

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	very weak to weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2521

KWS JOY

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium (3-5 per nerve)
Rachilla hair type	short
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

KWS MERIDIAN**CHARACTERS IN DETAIL****Ear**

Number of rows	six
Density	medium
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2879

KWS MOSELLE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax to medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect

WINTER BARLEY

AFP 2/2728

KWS ORWELL

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	long
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	weak
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	erect to semi-erect

Variety Name: **KWS Rose**

Crop:

Breeders Reference:

AFP No: 2/2730

Breeder:

Translated from:

Origin:

Date: 24 March 2015

Code: B2730

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Strong
Ear- attitude:	Semi - recurved to recurved
Ear- length:	Medium
*Ear- number of rows:	Two
*Ear- density:	Lax
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Very weak
*Awn- length compared to ear:	Short to medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	Parallel to weakly divergent
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	Medium
Collar type:	
Rachis- length of first segment:	Medium to long
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Very weak to weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

20 May 2015

WINTER BARLEY

AFP 2/2433

KWS SALSA

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium to long
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	weak to medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2580

KWS TONIC

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	dense
Length	medium
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	short to medium
Curvature of first segment	very weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	very weak to weak
Lemma nerves - at early dough	weak to medium
Awn tips	medium

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2725

KWS TOWER

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	erect to semi-erect

LG FLYNN**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	medium to strong
Lemma nerves - at early dough	medium to strong
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect

LG MOUNTAIN**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude	erect to semi-erect

WINTER BARLEY

AFP 2/2890

LIBRA

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	medium
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	50% equal:50% longer (segregation)

Grain

Kernel - colour of aleurone layer	50% coloured:50% whitish (segregation)
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	medium
Lemma nerves - at early dough	weak to medium
Awn tips	medium

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude	semi-erect to horizontal

WINTER BARLEY

AFP 2/2809

LG RAPTURE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

Variety Name: **Lucienne**

Crop:

Breeders Reference: BE2007011006

AFP No: 2/2885

Breeder: LS Plant Breeding Ltd

Translated from: German

Origin:

Date: 23 March 2017

Code: LUCI

*Plant - growth habit:	Semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Medium
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Absent or very weak
Ear- attitude:	Semi erect to horizontal
Ear- length:	Short
*Ear- number of rows:	More than two
*Ear- density:	Medium
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Medium to strong
*Awn- length compared to ear:	Long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	
Lemma nerves- anthocyanin colouration:	
*Plant height:	Long to very long
Collar type:	
Rachis- length of first segment:	Medium to long
Rachis- width of first segment:	
Rachis- curvature of first segment:	Weak to medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Weakly coloured
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Weak to medium
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

16 May 2018

MACKIE**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax to medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium to long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak
Ear - attitude	erect to semi-erect

WINTER BARLEY

AFP 2/2736

MALTESSE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium to long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

MARIS OTTER**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium to long
Shape	parallel

Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

Rachis

Length of first segment	medium to long
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	medium
Awn tips	medium

Plant growth characters

Plant - growth habit	semi-prostrate to prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late to very late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	horizontal

WINTER BARLEY

AFP 2/2452

MATROS

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax to medium
Length	medium to long
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	short to medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	very strong
Lemma nerves - at early dough	strong to very strong
Awn tips	strong to very strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	strong
Ear - attitude 21 days after ear emerg.	semi-recurved to recurved

WINTER BARLEY

AFP 2/2382

MEGANE

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	medium to dense
Length	medium
Shape	parallel

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium
Curvature of first segment	very weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	horizontal

WINTER BARLEY

AFP 2/2514

MEZMAAR

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	short to medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/1318

PEARL

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium (3-5 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	medium to strong
Lemma nerves - at early dough	very weak to weak
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	strong
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

WINTER BARLEY

AFP 2/2077

PELICAN

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	dense
Length	medium to long
Shape	

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	
Sterile spikelet - development	
Median spikelet - length of glume+awn cf grain	longer

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	weak
Awn tips	very weak to weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/2651

PERSEUS

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	short to medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	strong to very strong
Awn tips	strong

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2306

PURDEY

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium to long
Shape	parallel

Awn

Length compared to ear	shorter to +/- equal
------------------------	----------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	semi-prostrate to prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

WINTER BARLEY

AFP 2/2601

QUADRA

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	medium
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	short to medium
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	very weak to weak
Lemma nerves - at early dough	very weak to weak
Awn tips	weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2671

QUADRIGA

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	dense
Length	short to medium
Shape	parallel

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2065

RETRIEVER

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium to strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	medium
Lemma nerves - at early dough	strong
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

Variety Name: **Rubinesse**

Crop: Winter Barley

Breeders Reference: SC 82909 NH

AFP No:

Breeder: Secobra Recherches S A

Translated from: German

Origin:

Date: 28 March 2017

Code: RUBI

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Late
*Flag leaf- anthocyanin colouration of auricles:	Strong to very strong
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Medium to strong
*Ear- glaucosity:	Strong
Ear- attitude:	Semi erect to horizontal
Ear- length:	Medium
*Ear- number of rows:	Two
*Ear- density:	Medium to dense
Ear- shape:	Tapering to parallel
*Awn- anthocyanin colouration of the tips:	Very strong
*Awn- length compared to ear:	Short to medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	Divergent
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	Short
Collar type:	
Rachis- length of first segment:	Short to medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	`
Grain- spiculation of inner lateral nerves:	Weak to medium
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Strong
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

28 March 2017

Variety Name: **Sabine F**

Crop: Winter Barley

Breeders Reference:

AFP No: 2/2231

Breeder: Saat-zucht Filbeck

Translated from: German

Origin:

Date: 29 November 2007

Code: SABI

*Plant - growth habit:	intermediate
*Lower leaves - hairiness of leaf sheaths:	present
*Time of ear emergence:	late
*Flag leaf- anthocyanin colouration of auricles:	absent or very weak
Frequency of plants with recurved flag leaves:	high to very high
Flag leaf- glaucosity of sheath:	strong
*Ear- glaucosity:	weak to medium
Ear- attitude:	semi-erect to horizontal
Ear- length:	medium to long
*Ear- number of rows:	two
*Ear- density:	medium to dense
Ear- shape:	parallel
*Awn- anthocyanin colouration of the tips:	absent or very weak
*Awn- length compared to ear:	short to medium (shorter to +/- equal to ear)
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	parallel
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	medium
Collar type:	
Rachis- length of first segment:	long to very long
Rachis- width of first segment:	
Rachis- curvature of first segment:	weak to medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	whitish ("white")
*Grain- rachilla hair type:	long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	weak (1-2 per nerve)
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	absent or very weak
*Grain- ventral furrow - presence of hairs:	absent
*Grain- husk:	present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	clasping ("collar" type)
*Seasonal type:	winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

31 March 2015

SAFFRON**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	short to medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	absent or very weak
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

WINTER BARLEY

AFP 2/2228

SALLING

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	parallel

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

Variety Name: **SANDRA**

Crop:

Breeders Reference: BAUB 19104

AFP No:

Breeder: Berthold Bauer

Translated from: German

Origin:

Date: 02 April 2015

Code: SANDR

*Plant - growth habit:	Intermediate to semiprostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	High
Flag leaf- glaucosity of sheath:	Medium to strong
*Ear- glaucosity:	Weak to medium
Ear- attitude:	Horizontal to semi-recurved
Ear- length:	Medium
*Ear- number of rows:	Two
*Ear- density:	Medium
Ear- shape:	Tapering to parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	Parallel
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	
Collar type:	
Rachis- length of first segment:	Medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Weak
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Absent or very weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter

Extra characters:

* = Characteristics which should always be included in the description of the variety. 28 March 2017

WINTER BARLEY

AFP 2/1717

SEQUEL

CHARACTERS IN DETAIL

Ear

Number of rows	six
Density	medium
Length	short to medium
Shape	tapering to parallel

Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

Rachis

Length of first segment	short
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	
Sterile spikelet - development	
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

WINTER BARLEY

AFP 2/2742

SHADOW

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	short to medium
Shape	tapering

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	medium to long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/2091

SHANGRILA

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	dense to very dense
Length	short
Shape	

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	short
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	
Sterile spikelet - development	
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	very weak to weak
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	very early to early
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2454

SINATRA

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

Variety Name: **SOBELL**

Crop: Winter Barley

Breeders Reference: SJ 128113

AFP No:

Breeder: SEJET

Translated from: Danish

Origin:

Date: 15 May 2018

Code: SOBE

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium to late
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Medium to strong
*Ear- glaucosity:	Medium
Ear- attitude:	Horizontal to semi recurved
Ear- length:	Medium
*Ear- number of rows:	Two
*Ear- density:	Lax to medium
Ear- shape:	Tapering to parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	Non or rudimentary
Sterile spikelet-shape of tip:	NA
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	Absent or very weak
*Plant height:	
Collar type:	
Rachis- length of first segment:	Short to medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium to strong
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Absent or very weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

16 May 2018

WINTER BARLEY

AFP 2/2513

SOLOMAN

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	short
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/2822

SUNNINGDALE

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	medium
Length	short
Shape	parallel

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2745

SURGE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	short to medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	weak to medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2083

SUZUKA

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium
Shape	tapering to parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	short to medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early
Flag leaf - glaucosity of sheath	weak to medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

WINTER BARLEY

AFP 2/2444

SY BAMBOO

CHARACTERS IN DETAIL

Ear

Number of rows	more than two
Density	medium
Length	medium
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	longer

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate
Plant - length	long to very long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2888

SYCKLE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	strong
Awn tips	medium to strong

Plant growth characters

Plant - growth habit	intermediate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	strong
Ear - attitude	semi-erect

Variety Name: **SY TEPEE**

Crop:

Breeders Reference: SY 209-66

AFP No: 2/

Breeder: Sygenta Seeds Ltd

Translated from:

Origin: France

Date: 12 October 2012

Code: SYTEP

*Plant - growth habit:	Semi-prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Early
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	Medium to high
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Horizontal
Ear- length:	Medium to long
*Ear- number of rows:	Two
*Ear- density:	Medium
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Medium to long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	N/A
Sterile spikelet-shape of tip:	N/A
Sterile spikelet- length of lemma:	N/A
Median spikelet- length of glume + awn of grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	Medium
Collar type:	
Rachis- length of first segment:	Medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Medium to strong
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter type
Extra characters:	Ear - Development of sterile spiklets - Non or Rudimentary

* = Characteristics which should always be included in the description of the variety.

28 March 2018

WINTER BARLEY

AFP 2/2443

SY VENTURE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	very strong
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/2517

TALISMAN

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short (shorter than ear)
------------------------	--------------------------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	strong
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

WINTER BARLEY

AFP 2/2602

TETRIS

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	lax
Length	long
Shape	tapering

Awn

Length compared to ear	short (shorter than ear)
------------------------	--------------------------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	strong to very strong
Awn tips	strong

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

TOREROO**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	medium
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	medium
Lemma nerves - at early dough	medium
Awn tips	medium

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude	semi-erect to horizontal

WINTER BARLEY

AFP 2/2374

TRICK

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	parallel

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

WINTER BARLEY

AFP 2/2303

WINSOME

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	short
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	medium to strong
Awn tips	strong

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

WINTMALT

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering to parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	parallel to weakly divergent
Sterile spikelet - development	full
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	short
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

Variety Name: **Verity**

Crop: Winter Barley

Breeders Reference:

AFP No: 2/2750

Breeder:

Translated from: German

Origin:

Date: 01 April 2015

Code: VERI

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Very Strong
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Semi erect
Ear- length:	Short
*Ear- number of rows:	More than two
*Ear- density:	Lax to medium
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	N/A - Deficiens
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn of grain:	Longer
Lemma nerves- anthocyanin colouration:	
*Plant height:	Long
Collar type:	
Rachis- length of first segment:	Medium to long
Rachis- width of first segment:	
Rachis- curvature of first segment:	Very weak to weak
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Medium to strong
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

09 April 2015

WINTER BARLEY 2015



16/05/2018 V1

VOLUME**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	medium
Shape	parallel

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium
Curvature of first segment	absent or very weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	very weak to weak
Awn tips	weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

ZZOOM**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	medium
Length	short to medium
Shape	parallel

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	short to medium
Curvature of first segment	absent or very weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn of grain	longer

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	very strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude	semi-erect to horizontal